

The Picard rank conjecture for the Hurwitz spaces of degree up to five

Anand Deopurkar

Anand Patel

May 15, 2014

Abstract

We prove that the rational Picard group of the simple Hurwitz space $\mathcal{H}_{d,g}$ is trivial for d up to five. We also relate the rational Picard groups of the Hurwitz spaces to the rational Picard groups of the Severi varieties of nodal curves on Hirzebruch surfaces.

Keywords. Hurwitz space, Severi variety, Picard group

0 Introduction

Let $\mathcal{H}_{d,g}$ be the simple Hurwitz space which parametrizes isomorphism classes of simply branched degree d covers of genus zero curves by genus g curves. Although $\mathcal{H}_{d,g}$ has been studied classically, many fundamental questions about its geometry are still unanswered. The goal of this paper is to address one such question, the question of its Picard group. It is conjectured (for example, [DE96]) that the rational Picard group $\text{Pic}_{\mathbb{Q}}(\mathcal{H}_{d,g})$ is trivial. We call this the *Picard rank conjecture* for $\mathcal{H}_{d,g}$. Our main result is a proof of this conjecture for $d \leq 5$.

Theorem A. The rational Picard group of $\mathcal{H}_{d,g}$ is trivial for $d \leq 5$.

In the main text, Theorem A is divided into the case of degree 3 (Proposition 3.3), degree 4 (Proposition 4.10), and degree 5 (Proposition 5.4).

The Picard rank conjecture was known for $d = 2$ and 3. For $d = 2$, it was proved by Cornalba and Harris [CH88, Lemma 4.5], and for $d = 3$ by Stankova-Frenkel [SF00,

A. Deopurkar: Department of Mathematics, Columbia University, 2990 Broadway, New York, NY 10027; e-mail: anandrd@math.columbia.edu

A. Patel: Department of Mathematics, Boston College, Carney Hall, Chestnut Hill, MA 02467; e-mail: anand.patel@bc.edu

Mathematics Subject Classification (2010): Primary 14H10; Secondary 14H30

§ 12.2]. In these cases, now there are more refined results about the Picard group of the moduli stacks; see [Cor07] for $d = 2$ and [BV12] for $d = 3$.

The conjecture is also known for $d > 2g - 2$. In this range, the map $\mathcal{H}_{d,g} \rightarrow \mathcal{M}_g$ is a fibration, where \mathcal{M}_g is the moduli space of smooth curves of genus g . An analysis of this fibration shows that $\text{Pic}_{\mathbf{Q}}(\mathcal{H}_{d,g}) = 0$ if and only if $\text{Pic}_{\mathbf{Q}}(\mathcal{M}_g) \cong \mathbf{Q}$ (see, for example, [Moc95] or [DE96, § 3]). Thus, the conjecture for $d > 2g - 2$ follows from Harer's theorem [Har83].

We briefly explain the rationale behind the conjecture. Let us blur the distinction between the coarse moduli spaces and the fine moduli stacks. This is harmless, since we are concerned with the rational Picard group. Let us also take $d \geq 4$ (the discussion holds for $d = 2, 3$ with minor modifications). Denote by $\tilde{\mathcal{H}}_{d,g}$ the partial compactification of $\mathcal{H}_{d,g}$ that parametrizes covers $[\alpha: C \rightarrow \mathbf{P}^1]$ where C is allowed to be nodal, but still irreducible, and α need not be simply branched. Let $\alpha: \mathcal{C} \rightarrow \mathcal{P}$ be the universal family over $\tilde{\mathcal{H}}_{d,g}$, where $\rho: \mathcal{C} \rightarrow \tilde{\mathcal{H}}_{d,g}$ is a family of irreducible, at worst nodal curves of arithmetic genus g , and $\pi: \mathcal{P} \rightarrow \tilde{\mathcal{H}}_{d,g}$ a family of smooth curves of genus 0. From this data, we can construct three ‘tautological’ divisor classes on $\tilde{\mathcal{H}}_{d,g}$ given by

$$\rho_*(c_1(\omega_\rho)^2), \quad \rho_*(c_1(\omega_\rho)\alpha^*c_1(\omega_\pi)), \quad \text{and} \quad \rho_*([\delta_\rho]).$$

Here ω stands for the relative dualizing sheaf and δ for the singular locus. It is easy to check that the three tautological classes are \mathbf{Q} -linearly independent. On the other hand, $\tilde{\mathcal{H}}_{d,g} \setminus \mathcal{H}_{d,g}$ is a union of three irreducible divisors, namely the locus Δ where C is singular, the locus T where α has a higher order ramification point, and the locus D where α has two ramification points over a branch point. It is also easy to check that the classes of Δ , T , and D are \mathbf{Q} -linearly independent. Thus, $\text{Pic}_{\mathbf{Q}}(\mathcal{H}_{d,g}) = 0$ is equivalent to $\text{Pic}_{\mathbf{Q}}(\tilde{\mathcal{H}}_{d,g})$ being generated by the tautological classes. The Picard rank conjecture thus expresses the often-satisfied expectation that there are no other divisor classes than the tautological ones.

We now outline our strategy for proving Theorem A. Let $\alpha: C \rightarrow \mathbf{P}^1$ be a degree d cover. Then C embeds in a \mathbf{P}^{d-2} -bundle over \mathbf{P}^1 , which we denote by $\mathbf{P}E \rightarrow \mathbf{P}^1$. Thanks to the work of Casnati and Ekedahl, the resolution of the ideal of C in $\mathbf{P}E$ can be described explicitly. The terms in this resolution involve (twists of) vector bundles on \mathbf{P}^1 [CE96]. Let $U \subset \tilde{\mathcal{H}}_{d,g}$ be the open locus where these vector bundles are the most generic. The key steps in our proof are the following.

1. Identify the divisorial components of $\tilde{\mathcal{H}}_{d,g} \setminus U$.
2. Express U as a (successive) quotient of an open subset of an affine space by actions of linear algebraic groups.
3. Use the previous two steps to get a bound on the Picard rank of $\tilde{\mathcal{H}}_{d,g}$, and in turn, the Picard rank of $\mathcal{H}_{d,g}$.

Needless to say, we are able to carry out all three steps only for $d \leq 5$. However, we can carry out parts of step (1) in general. For step (2), we highlight that it remains unknown in general whether one can dominate $\tilde{\mathcal{H}}_{d,g}$ by an affine space for $d \geq 6$.

To analyze $\tilde{\mathcal{H}}_{d,g} \setminus U$, we must analyze the loci in $\tilde{\mathcal{H}}_{d,g}$ where the bundle E and the vector bundles appearing in the resolution of C are unbalanced. We call these loci the Maroni loci and the Casnati–Ekedahl loci, respectively. We spend significant effort on understanding the decomposition of $\tilde{\mathcal{H}}_{d,g}$ into these loci. Contained in Section 2, the results of this analysis may be of independent interest.

A key tool in our analysis is a construction that relates the Maroni loci to the Severi varieties of Hirzebruch surfaces. Originally due to Ohbuchi [Ohb97], this ‘associated scroll construction’ allows us to get the required dimension estimates. The key input here is a theorem of Tyomkin that guarantees that the Severi varieties of Hirzebruch surfaces are irreducible of the expected dimension [Tyo07].

The associated scroll construction also lets us relate the Picard ranks of the Hurwitz spaces to the Picard ranks of the Severi varieties. To state our result, let us denote by $\mathcal{U}_g(\mathbf{F}_m, d\tau)$ the space of irreducible nodal curves of geometric genus g in the linear system $|d\tau|$ on the Hirzebruch surface \mathbf{F}_m , where τ is the section with self-intersection m .

Theorem B. Let $m \geq \lfloor (g + d - 1)/(d - 1) \rfloor$. Then $\text{Pic}_{\mathbf{Q}} \mathcal{U}_g(\mathbf{F}_m, d\tau) = 0$ implies $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{d,g} = 0$.

Let $m \geq \lceil 2(g + d - 1)/(d - 1) \rceil$. Then $\text{Pic}_{\mathbf{Q}} \mathcal{U}_g(\mathbf{F}_m, d\tau) = 0$ if and only if $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{d,g} = 0$.

In the main text, Theorem B is Theorem 6.7.

0.1 Notation

We work with a few different versions of the Hurwitz spaces. We assemble their definitions here. We work over the field \mathbf{C} of complex numbers. By a curve, we mean a connected, proper, reduced scheme of finite type over \mathbf{C} . Throughout, assume that $g \geq 3$.

$\mathcal{H}_{d,g}$: This is the coarse moduli space of $[\alpha: C \rightarrow \mathbf{P}^1]$, where C is a smooth curve of genus g and α a finite map of degree d with simple branching (that is, the branch divisor of α is supported at $2g + 2d - 2$ distinct points). Two such covers $[\alpha_1: C_1 \rightarrow \mathbf{P}^1]$ and $[\alpha_2: C_2 \rightarrow \mathbf{P}^1]$ are considered isomorphic if there are isomorphisms $\phi: C_1 \rightarrow C_2$ and $\psi: \mathbf{P}^1 \rightarrow \mathbf{P}^1$ such that $\alpha_2 \circ \phi = \psi \circ \alpha_1$.

$\tilde{\mathcal{H}}_{d,g}$: This is the coarse moduli space of $[\alpha: C \rightarrow \mathbf{P}^1]$, where C is an irreducible, at worst nodal curve of arithmetic genus g , and α a finite map of degree d . The isomorphism condition is the same as that for $\mathcal{H}_{d,g}$.

$\mathcal{H}_{d,g}^\dagger$: This is like $\mathcal{H}_{d,g}$, but with ‘framed’ target \mathbf{P}^1 . The objects it parametrizes are $[\alpha: C \rightarrow \mathbf{P}^1]$ as in the description of $\mathcal{H}_{d,g}$, but $[\alpha_1: C_1 \rightarrow \mathbf{P}^1]$ and $[\alpha_2: C_2 \rightarrow \mathbf{P}^1]$ are considered isomorphic if there is an isomorphism $\phi: C_1 \rightarrow C_2$ such that $\alpha_2 \circ \phi = \alpha_1$.

$\tilde{\mathcal{H}}_{d,g}^\dagger$: This is like $\tilde{\mathcal{H}}_{d,g}$, but with framed target \mathbf{P}^1 .

All four are irreducible quasi-projective varieties with at worst quotient singularities. In particular, they are normal and \mathbf{Q} -factorial. The group $\text{Aut } \mathbf{P}^1 = \text{PGL}_2$ acts on the framed versions. The unframed versions are the quotients by this action in the sense that the fibers of the morphism from the framed space to the unframed space are precisely the PGL_2 orbits. We have

$$\dim \mathcal{H}_{d,g} = \dim \tilde{\mathcal{H}}_{d,g} = 2g + 2d - 5,$$

and

$$\dim \mathcal{H}_{d,g}^\dagger = \dim \tilde{\mathcal{H}}_{d,g}^\dagger = 2g + 2d - 2.$$

In addition, we work with the following Severi varieties:

$\mathcal{U}_g(\mathbf{F}_m, d\tau)$: This is the locus of irreducible nodal curves of geometric genus g in the linear series $|d\tau|$ in the Hirzebruch surface \mathbf{F}_m . Here $\tau \subset \mathbf{F}_m$ is the section of self-intersection m .

$\mathcal{V}_g(\mathbf{F}_m, d\tau)$: This is the closure of $\mathcal{U}_g(\mathbf{F}_m, d\tau)$ in the projective space $|d\tau|$.

$\mathcal{V}_g^{\text{irr}}(\mathbf{F}_m, d\tau)$: This is the open subset of reduced and irreducible curves in $\mathcal{V}_g(\mathbf{F}_m, d\tau)$.

We do not distinguish between a vector bundle and the corresponding locally free sheaf. Note that the vector bundle associated to the locally free sheaf F is the relative Spec of the symmetric algebra on F^\vee .

1 Preliminaries

In this expository section, we recall two key results. The first describes the Picard group of the quotient of a variety by a group action. The second is a structure theorem for finite covers which enables us to describe a large open subset of the Hurwitz space as such a quotient.

1.1 Picard groups of quotients

Let G be a linear algebraic group acting on a variety X . Denote by $\text{Pic}_G X$ the group of G -linearized line bundles on X . Forgetting the G -linearization gives a homomorphism $\text{Pic}_G X \rightarrow \text{Pic } X$.

Proposition 1.1. [KKV89, Lemma 2.2 + Proposition 2.3] *For a connected linear algebraic group G acting on an irreducible variety X , we have an exact sequence*

$$\chi(G) \rightarrow \mathrm{Pic}_G X \rightarrow \mathrm{Pic} X,$$

where $\chi(G)$ is the group of (algebraic) characters of G . Furthermore, if X is normal, then the sequence has an extension by a homomorphism $\mathrm{Pic} X \rightarrow \mathrm{Pic} G$.

Let $\pi: X \rightarrow Y$ be a morphism that is equivariant with the trivial G action on Y . Let L be a line bundle on Y . The pullback π^*L carries a natural G -linearization. We thus have a homomorphism $\mathrm{Pic} Y \rightarrow \mathrm{Pic}_G X$.

Proposition 1.2. *Let X and Y be irreducible normal varieties, G a linear algebraic group acting on X , and $\pi: X \rightarrow Y$ a surjective morphism, equivariant with the trivial action on Y . Suppose the fibers of π consist of single G -orbits. Then the map $\mathrm{Pic} Y \rightarrow \mathrm{Pic}_G X$ is injective and we have*

$$\mathrm{rk} \mathrm{Pic} Y \leq \mathrm{rk} \chi(G) + \mathrm{rk} \mathrm{Pic} X.$$

Furthermore, if G is reductive and the stabilizers G_x are finite, then we have an isomorphism

$$\mathrm{Pic} Y \otimes \mathbf{Q} \xrightarrow{\sim} \mathrm{Pic}_G X \otimes \mathbf{Q}.$$

Proof. Suppose L is a line bundle on Y such that π^*L is trivial as a G -linearized line bundle. Then π^*L has a G -invariant nowhere-vanishing section. We claim that such a section descends to a nowhere-vanishing section of L on Y . The crucial point is that in our setup, Y is a geometric quotient of X [MFK94, Proposition 0.2]. That is, for every open $U \subset Y$, the preimage $\pi^{-1}U$ is open and the functions on U are the invariant functions on $\pi^{-1}U$:

$$\Gamma(U, O_Y) = \Gamma(\pi^{-1}U, O_X)^G.$$

It follows that the sections of L on U are the invariant sections of π^*L on $\pi^{-1}(U)$:

$$\Gamma(U, L) = \Gamma(\pi^{-1}U, \pi^*L)^G.$$

Thus, a G -invariant section σ of π^*L on X gives a section $\bar{\sigma}$ of L on Y . It is easy to check that if σ is nowhere-vanishing, so is $\bar{\sigma}$.

The bound on $\mathrm{rk} \mathrm{Pic} Y$ follows from the injectivity and Proposition 1.1. For the last statement, we use the characterization of the image of $\mathrm{Pic} Y \rightarrow \mathrm{Pic}_G X$ from [KKV89, Proposition 4.2]: a G -linearized line bundle L is in the image if and only if for every $x \in X$, the stabilizer group G_x acts trivially on the fiber L_x . Since the stabilizers are finite, we can arrange this by passing to a large enough power of L . \square

We end with a simple application.

Proposition 1.3. *Let $U \subset \widetilde{\mathcal{H}}_{d,g}$ be any open subset and U^\dagger its preimage under $\widetilde{\mathcal{H}}_{d,g}^\dagger \rightarrow \widetilde{\mathcal{H}}_{d,g}$. Then*

$$\mathrm{rk} \mathrm{Pic} U = \mathrm{rk} \mathrm{Pic} U^\dagger.$$

Proof. Apply Proposition 1.1 and Proposition 1.2 with $G = \mathrm{PGL}_2$, $X = U^\dagger$, and $Y = U$. \square

1.2 The Casnati–Ekedahl structure theorem

Let X and Y be integral schemes and $\alpha: X \rightarrow Y$ a finite flat Gorenstein morphism of degree $d \geq 3$. The map α gives an exact sequence

$$0 \rightarrow O_Y \rightarrow \alpha_* O_X \rightarrow E_\alpha^\vee \rightarrow 0, \quad (1.1)$$

where $E = E_\alpha$ is a vector bundle of rank $(d-1)$ on Y , called the *Tschirnhausen bundle* of α . Denote by ω_α the dualizing sheaf of α . Applying $\mathrm{Hom}_Y(-, O_Y)$ to (1.1), we get

$$0 \rightarrow E \rightarrow \alpha_* \omega_\alpha \rightarrow O_Y \rightarrow 0. \quad (1.2)$$

The map $E \rightarrow \alpha_* \omega_\alpha$ induces a map $\alpha^* E \rightarrow \omega_\alpha$.

Theorem 1.4. *[CE96, Theorem 2.1] In the above setup, $\alpha^* E \rightarrow \omega_\alpha$ gives an embedding $\iota: X \rightarrow \mathbf{P}E$ with $\alpha = \pi \circ \iota$, where $\pi: \mathbf{P}E \rightarrow Y$ is the projection. Moreover, the subscheme $X \subset \mathbf{P}E$ can be described as follows.*

1. *The resolution of O_X as an $O_{\mathbf{P}E}$ module has the form*

$$\begin{aligned} 0 \rightarrow \pi^* N_{d-2}(-d) \rightarrow \pi^* N_{d-3}(-d+2) \rightarrow \pi^* N_{d-4}(-d+3) \rightarrow \dots \\ \dots \rightarrow \pi^* N_2(-3) \rightarrow \pi^* N_1(-2) \rightarrow O_{\mathbf{P}E} \rightarrow O_X \rightarrow 0, \end{aligned} \quad (1.3)$$

where the N_i are vector bundles on Y . Restricted to a point $y \in Y$, this sequence is the minimal free resolution of $X_y \subset \mathbf{P}E_y$.

2. *The ranks of the N_i are given by*

$$\mathrm{rk} N_i = \frac{i(d-2-i)}{d-1} \binom{d}{i+1},$$

3. *We have $N_{d-2} \cong \pi^* \det E$. Furthermore, the resolution is symmetric, that is, isomorphic to the resolution obtained by applying $\mathrm{Hom}_{O_{\mathbf{P}E}}(-, N_{d-2}(-d))$.*

The branch divisor of $\alpha: X \rightarrow Y$ is given by a section of $(\det E)^{\otimes 2}$. In particular, if X is a curve of (arithmetic) genus g , α has degree d , and $Y = \mathbf{P}^1$, then

$$\mathrm{rk} E = d-1 \text{ and } \deg E = g + d - 1. \quad (1.4)$$

2 The Maroni and Casnati–Ekedahl loci

Consider a cover $\alpha: C \rightarrow \mathbf{P}^1$ and the relative canonical embedding $C \subset \mathbf{P}E_\alpha$. Since vector bundles on \mathbf{P}^1 split as direct sums of line bundles, the vector bundle E_α , and the higher syzygy bundles N_i appearing in Theorem 1.4 are discrete invariants of α . We thus get a decomposition of the Hurwitz space into locally closed subsets where the isomorphism type of the bundles E_α and N_i are constant. This section is devoted to the analysis of some of these locally closed subvarieties, particularly their dimensions. We only consider the bundle E_α and $F_\alpha := N_1$. Note that

$$E_\alpha = \ker(\alpha_*\omega_\alpha \rightarrow O_Y) \text{ and } F_\alpha = \alpha_*I_C(2),$$

where $I_C \subset O_{\mathbf{P}E_\alpha}$ is the ideal sheaf of C .

Definition 2.1. For vector bundles E and F on \mathbf{P}^1 , define the following closed subvarieties of $\mathcal{H}_{d,g}^\dagger$:

$$\begin{aligned} M(E, F) &:= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid E_\alpha \cong E \text{ and } F_\alpha \cong F\}}, \\ M(E) &:= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid E_\alpha \cong E\}}, \\ C(F) &:= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid F_\alpha \cong F\}}. \end{aligned}$$

Call $M(E)$ the *Maroni loci* and $C(F)$ the *Casnati–Ekedahl loci*. Define subvarieties $\widetilde{M}(E, F)$, $\widetilde{M}(E)$, and $\widetilde{C}(F)$ of $\mathcal{H}_{d,g}^\dagger$ analogously.

Abusing notation, we denote the images of these loci in the unframed versions $\mathcal{H}_{d,g}$ and $\widetilde{\mathcal{H}}_{d,g}$ by the same letters. The framed versus unframed setting is usually clear by context, and sometimes irrelevant, for example in discussing the codimensions. We caution the reader that these loci are not necessarily irreducible or of expected dimension (Example 4.3, Example 4.4). Even determining whether they are non-empty remains a challenge in full generality.

2.1 The associated scroll construction

To analyze the Maroni loci $M(E)$, we associate to a cover of \mathbf{P}^1 a curve on a Hirzebruch surface. The construction is originally due to Ohbuchi [Ohb97]. Let C be an irreducible curve of arithmetic genus g and $\alpha: C \rightarrow \mathbf{P}^1$ a finite cover of degree d . Let ζ be a global section of $O_C(m) = \alpha^*O_{\mathbf{P}^1}(m)$ that projects to a nonzero section of $E_\alpha^\vee(m)$. In other words, ζ is not a pullback of a section from \mathbf{P}^1 . The section ζ gives a map from C to the total space of the line bundle $O(m)$ over \mathbf{P}^1 . Let $\mathbf{F}_m = \text{Proj}(O \oplus O(-m))$ be the

Hirzebruch surface that compactifies this total space. We thus get the diagram

$$\begin{array}{ccc} C & \xrightarrow{\nu} & \mathbf{F}_m \\ & \searrow \alpha & \downarrow \pi \\ & & \mathbf{P}^1 \end{array}$$

Let $\sigma \subset \mathbf{F}_m$ be the directrix and $\tau \subset \mathbf{F}_m$ the section disjoint from σ (so that $\sigma^2 = -m$ and $\tau^2 = m$). By construction, $\nu(C) \subset \mathbf{F}_m$ avoids the directrix σ . Suppose C is smooth and $\alpha: C \rightarrow \mathbf{P}^1$ does not factor nontrivially. Then ν is birational onto its image, and therefore $\nu(C)$ is a reduced and irreducible element of the linear system $|d\tau|$. By the following proposition, $\nu(C)$ is a point in the Severi variety $\mathcal{V}_g(\mathbf{F}_m, d\tau)$.

Proposition 2.2. *A reduced and irreducible curve on \mathbf{F}_m of geometric genus g in the linear system $|d\tau|$ is a flat limit of irreducible nodal curves of geometric genus g .*

Proof. Let $\bar{C} \subset \mathbf{F}_m$ be such a reduced and irreducible curve. Let $C \rightarrow \bar{C}$ the normalization. Denote by ν the composite map $\nu: C \rightarrow \mathbf{F}_m$. Let \mathcal{M} be a component of the Kontsevich space of maps $\mathcal{M}_g(\mathbf{F}_m, d\tau)$ containing ν . Let N_ν be the normal sheaf of ν ; this is the cokernel of $T_C \rightarrow \nu^*T_{\mathbf{F}_m}$. Then, we have a lower bound: $\dim \mathcal{M} \geq \chi(N_\nu)$. Since

$$\chi(N_\nu) = \chi(\nu^*T_{\mathbf{F}_m}) - \chi(T_C) = g - \deg(K_{\mathbf{F}_m} \cdot \bar{C}) - 1,$$

we get

$$\dim \mathcal{M} \geq g - \deg(K_{\mathbf{F}_m} \cdot \bar{C}) - 1.$$

By [Har86, Proposition 2.2], a general $\nu_{\text{gen}}: C_{\text{gen}} \rightarrow \mathbf{F}_m$ in \mathcal{M} is birational onto its image and the image has only nodes as singularities. \square

We can make the construction in a family. Let M be a reduced scheme, $\rho: C \rightarrow M$ a generically smooth family of reduced and irreducible curves of genus g , and $\alpha: C \rightarrow \mathbf{P}^1 \times M$ a finite flat M -morphism of degree d . Set $O_C(m) = \alpha^*O(m)$. Assume that none of the fibers $\alpha_t: C_t \rightarrow \mathbf{P}^1$ factor nontrivially and $H^0(C_t, O_{C_t}(m))$ has constant rank. Then $\rho_*O_C(m)$ is a vector bundle on M . The trivial subbundle $H^0(\mathbf{P}^1, O(m)) \otimes_{O_M}$ maps injectively to $\rho_*O_C(m)$. Let U be the complement of the image of this map in the total space of $\rho_*O_C(m)$. Fiberwise, the sections of U correspond to the sections ζ which project nontrivially onto $E_\alpha^\vee(m)$. Then the associated scroll construction gives a morphism

$$U \rightarrow \mathcal{V}_g(\mathbf{F}_m, d\tau).$$

We will use this construction where M is a Maroni locus. As described, the construction depends on the existence of a universal family, and thus gives a morphism from the fine

moduli stack. But since $\mathcal{V}_g(\mathbf{F}_m, d\tau)$ is a scheme, we get a canonical induced map from the coarse space.

The following crucial result makes the above construction useful.

Theorem 2.3 ([Tyo07]). *All Severi varieties parametrizing irreducible curves on Hirzebruch surfaces are irreducible and of expected dimension. In particular, the variety $\mathcal{V}_g(\mathbf{F}_m, d\tau)$ is irreducible of dimension $dm + 2d + g - 1$.*

We also need the following result, which we prove for the lack of a reference.

Proposition 2.4. *Let $\bar{C} \subset \mathbf{F}_m$ be a general point of $\mathcal{V}_g(\mathbf{F}_m, d\tau)$ and $C \rightarrow \bar{C}$ the normalization. Then the composite $C \rightarrow \mathbf{P}^1$ is simply branched.*

Proof. In light of Theorem 2.3, it suffices to exhibit a particular \bar{C} of geometric genus g in $\mathcal{V}_g(\mathbf{F}_m, d\tau)$ whose normalization is simply branched over \mathbf{P}^1 . One way is to start with $X = \mathbf{P}^1$ and $\alpha: X \rightarrow \mathbf{P}^1$ a simply branched cover of degree d . Then $E_\alpha = O(1)^{\oplus(d-1)}$. Choosing a general section of $E_\alpha^\vee(m)$ gives $\nu: X \rightarrow \mathbf{F}_m$ such that $\nu(X)$ is nodal. It is easy to see that $\nu(X)$ is in the closure of $\mathcal{V}_g(\mathbf{F}_m, d\tau)$. Indeed, since the set of nodes of $\nu(X)$ impose independent conditions on $|K_{\mathbf{F}_m} + d\tau|$, they automatically impose independent conditions on $|d\tau|$ as well, and hence we may smooth out the required number of nodes of $\nu(X)$ to deform to a curve of geometric genus g . A general fiber of such a smoothing is the required \bar{C} . \square

Remark 2.5. We can realize the associated scroll construction geometrically as follows. The choice of a general global section ζ of $O_C(m)$ can be thought of as a choice of a geometric section $\sigma: \mathbf{P}^1 \rightarrow \mathbf{P}E$. In the \mathbf{P}^{d-2} fibers of $\pi: \mathbf{P}E \rightarrow \mathbf{P}^1$, we now have $d+1$ points: d points coming from the fibers of the map $\alpha: C \rightarrow \mathbf{P}^1$, and one more point provided by the section σ . For general $t \in \mathbf{P}^1$, these $d+1$ points will be in general position, and so will define a unique rational normal curve $R_t \subset \mathbf{P}E$. Consider the birationally ruled surface $S \subset \mathbf{P}E$ defined as the closure of the union of the R_t 's. S contains both σ and C , and is fibered over \mathbf{P}^1 . We contract all components of the fibers of the projection $\pi: S \rightarrow \mathbf{P}^1$ which do not meet the directrix σ . The resulting surface is \mathbf{F}_m , with σ being the directrix. The image of C under the contraction $S \rightarrow \mathbf{F}_m$ is the associated scroll construction.

For a vector bundle $E = O(a_1) \oplus \cdots \oplus O(a_n)$ on \mathbf{P}^1 , set

$$\lfloor E \rfloor = \min\{a_i\} \text{ and } \lceil E \rceil = \max\{a_i\}.$$

Given a cover $\alpha: C \rightarrow \mathbf{P}^1$, the associated scroll construction $\nu: C \rightarrow \mathbf{F}_m$ can be made for $m \geq \lfloor E_\alpha \rfloor$. Conversely, given a point $\bar{C} \in \mathcal{V}_g^{\text{irr}}(\mathbf{F}_m, d\tau)$, let $C \rightarrow \bar{C}$ be the normalization. Then the induced cover $\alpha: C \rightarrow \mathbf{P}^1$ has $\lfloor E_\alpha \rfloor \leq m$.

Proposition 2.6. *If $\widetilde{M}(E)$ is nonempty, then*

$$\lceil E \rceil \leq \frac{2g + 2d - 2}{d}. \quad (2.1)$$

Furthermore, if E_α comes from a cover $[\alpha: C \rightarrow \mathbf{P}^1]$, with C irreducible, and where α does not factor nontrivially, then

$$\frac{g + d - 1}{\binom{d}{2}} \leq \lfloor E_\alpha \rfloor \leq \frac{g + d - 1}{d - 1}. \quad (2.2)$$

Proof. The resolution of O_C in Theorem 1.4 tells us that $C \subset \mathbf{P}E_\alpha$ is not contained in any hyperplane divisor. Let h denote the hyperplane divisor class associated to $O_{\mathbf{P}E_\alpha}(1)$, and let f denote the class of the fiber of $\pi: \mathbf{P}E \rightarrow \mathbf{P}^1$. Set $N := \lceil E_\alpha \rceil$. Then the divisor class $h - Nf$ is effective. Since C is irreducible and does not lie in $(h - Nf)$, it intersects $(h - Nf)$ non-negatively. Since $h \cdot [C] = 2g + 2d - 2$, and $f \cdot [C] = d$, we conclude that $N \leq \frac{2d+2g-2}{d}$.

For the second inequality, we appeal to the associated scroll construction. Let $n := \lfloor E_\alpha \rfloor$. Since α does not factor, $\nu: C \rightarrow \mathbf{F}_n$ must be birational onto its image. Adjunction on \mathbf{F}_n gives

$$p_a(\nu(C)) = \binom{d}{2}n - (d - 1).$$

The second statement now follows from the inequality $g \leq p_a(\nu(C))$. \square

The following theorem of Ohbuchi [Ohb97] places a strong restriction on a large class of Tschirnhausen bundles E .

Proposition 2.7 ([Ohb97]). *Let $\alpha: C \rightarrow \mathbf{P}^1$ be a cover of degree d , with C irreducible, and where α does not factor nontrivially. Write $E_\alpha = O(a_1) \oplus \cdots \oplus O(a_{d-1})$ where $\lfloor E_\alpha \rfloor = a_1 \leq a_2 \leq \cdots \leq a_{d-1} = \lceil E_\alpha \rceil$. Then*

$$a_{i+1} - a_i \leq \lfloor E_\alpha \rfloor \quad \text{for } 1 \leq i \leq d - 2. \quad (2.3)$$

Remark 2.8. Proposition 2.7 implies the second inequality in Proposition 2.6.

Definition 2.9. We call a vector bundle E on \mathbf{P}^1 of rank $d - 1$ and degree $g + d - 1$ *tame* if it satisfies the inequalities (2.1), (2.2), and (2.3).

Notice that Proposition 2.6 and Proposition 2.7 imply that E_α is tame in the following two cases: α is simply branched, or d is prime. Indeed, in either case, the cover cannot factor non-trivially.

Denote by \rightsquigarrow the partial order on vector bundles on \mathbf{P}^1 given by $E \rightsquigarrow E'$ if E specializes to E' in a flat family. Define the finite set $\mathcal{T}[m]$ by

$$\mathcal{T}[m] := \{\text{Isomorphism classes of tame bundles } E \text{ of rank } d - 1, \text{ degree } g + d - 1, \text{ and } \lfloor E \rfloor = m\}.$$

Observe that $\mathcal{T}[m]$ contains an element $E[m]$ such that $E[m] \rightsquigarrow E$ for all $E \in \mathcal{T}[m]$. In other words, $E[m]$ is the most generic among all the bundles in $\mathcal{T}[m]$.

Theorem 2.10. *Let m be an integer satisfying $\frac{g+d-1}{\binom{d}{2}} \leq m \leq \frac{g+d-1}{d-1}$.*

1. *If $M(E)$ is nonempty, then E is a tame bundle.*
2. *If $\lfloor E \rfloor \leq m$ then $M(E) \subset M(E[m])$.*
3. *$M(E[m]) \subset M(E[m+1])$ for all m .*
4. *$M(E[m])$ is an irreducible subvariety of $\mathcal{H}_{d,g}^\dagger$ of codimension $g - (d-1)m + 1$ unless $m = \lfloor \frac{g+d-1}{d-1} \rfloor$, in which case $M(E[m]) = \mathcal{H}_{d,g}^\dagger$.*
5. *If d is prime, then all the statements above hold with $M(-)$ replaced by $\widetilde{M}(-)$ and $\mathcal{H}_{d,g}^\dagger$ replaced by $\widetilde{\mathcal{H}}_{d,g}^\dagger$.*

In the proof, we use a theorem of Coppens, which we state using our setup.

Theorem 2.11 ([Cop99]). *For all m satisfying $\frac{g+d-1}{\binom{d}{2}} \leq m \leq \frac{g+d-1}{d-1}$, there is a genus g and degree d cover $C \rightarrow \mathbf{P}^1$ with Tschirnhausen bundle $E[m]$. Moreover, C is birational onto its image under the associated scroll construction $C \rightarrow \mathbf{F}_m$.*

Proof of Theorem 2.10. We repeatedly use simultaneous normalization in the following way: Suppose we have a family $\mathcal{C} \rightarrow \Delta$ of reduced irreducible curves of geometric genus g . Then the normalization \mathcal{C}^ν of \mathcal{C} gives a family $\mathcal{C}^\nu \rightarrow \Delta$ of smooth curves of genus g [Tei80]. For the ease of reading, we do not make this process explicit every time.

The first statement follows from Proposition 2.6 and Proposition 2.7.

For the second statement, first note that if $[\overline{C}] \in \mathcal{V}_g(\mathbf{F}_m, d\tau)$ is a general point and $\nu: C \rightarrow \overline{C}$ the normalization, then $C \rightarrow \mathbf{P}^1$ is simply branched and has Tschirnhausen bundle $E[m]$. Indeed, we can get a $[\overline{C}] \in \mathcal{V}_g(\mathbf{F}_m, d\tau)$ with Tschirnhausen module $E[m]$ by applying the associated scroll construction to a cover given by Theorem 2.11. By Proposition 2.4, we may deform such $[\overline{C}]$ so that the normalization is simply branched. By the genericity of $E[m]$, the normalization of the deformed curve also has Tschirnhausen bundle $E[m]$. Now, suppose $\lfloor E \rfloor \leq m$ and $[C \rightarrow \mathbf{P}^1]$ is a point with Tschirnhausen bundle E . Then the associated scroll construction gives $\nu: C \rightarrow \mathbf{F}_m$. Since α is simply branched, ν is birational onto its image. Then $\nu(C)$ is the limit of curves in $\mathcal{V}_g(\mathbf{F}_m, d\tau)$ whose normalization has Tschirnhausen bundle $E[m]$. The second statement follows.

The third statement is a corollary of the second statement.

For the fourth statement, suppose $m = \lfloor \frac{g+d-1}{d-1} \rfloor$. Then $E[m]$ is balanced, so $M(E[m]) = \mathcal{H}_{d,g}^\dagger$. Suppose $m < \lfloor \frac{g+d-1}{d-1} \rfloor$. Let $U \subset \mathcal{V}_g(\mathbf{F}_m, d\tau)$ be the locus of nodal curves of geometric genus g whose normalization is simply branched over \mathbf{P}^1 . Then U is a smooth

open subset of $\mathcal{V}_g(\mathbf{F}_m, d\tau)$. Normalization of the universal family of curves in \mathbf{F}_m of geometric genus g gives a family of smooth curves of genus g with a simply branched map of degree d to \mathbf{P}^1 (induced from $\mathbf{F}_m \rightarrow \mathbf{P}^1$.) By definition, the image is in $M(E[m])$. We thus get a dominant map

$$q: U \rightarrow M(E[m]).$$

The fiber of q over $[\alpha: C \rightarrow \mathbf{P}^1]$ corresponds to the global sections of $O_C(m)$ that project non-trivially onto $E^\vee(m)$. For general $\alpha \in M(E[m])$, we have $E_\alpha = E[m]$. Also, since $m < \lfloor \frac{g+d-1}{d-1} \rfloor$, the bundle $E[m]$ has a unique $O(m)$ summand and all other summands have degree greater than m . Therefore, the general fiber of q has dimension $m + 2$. From the dimension of $\mathcal{V}_g(\mathbf{F}_m, d\tau)$, we get

$$\dim M(E[m]) = \dim \mathcal{V}_g(\mathbf{F}_m, d\tau) - (m + 2) = (d - 1)m + g + 2d - 3.$$

Since $\dim \mathcal{H}_{d,g}^\dagger = 2g + 2d - 2$, the fourth statement follows.

For the last statement, note that all the arguments hold for $\widetilde{M}(E)$ if d is prime, since the associated scroll construction $\nu: C \rightarrow \mathbf{F}_m$ is automatically birational onto its image. \square

Theorem 2.10 gives us good control on the dimensions of the Maroni loci for E based on the minimal summand of E . We must now consider those E which are non-generic, but nonetheless have the same minimal summand as the generic Tschirnhausen bundle. Set $k = \lfloor \frac{g+d-1}{d-1} \rfloor$. Then

$$E[k] = O(k)^{\oplus r} \oplus O(k+1)^{\oplus d-r-1},$$

where $0 < r \leq d - 1$. A general cover $\alpha \in \mathcal{H}_{d,g}^\dagger$ has $E[k]$ as its Tschirnhausen bundle. Let E' be any tame bundle, and set $s := h^0(E'^\vee(k))$. Upper semicontinuity implies $s \geq r$. Suppose $s > r$. Define

$$M^\circ(E') = \left\{ \alpha \in \mathcal{H}_{d,g}^\dagger \mid E_\alpha \cong E' \right\}.$$

Then $M^\circ(E')$ is locally closed, and $\overline{M^\circ(E')} = M(E')$.

Lemma 2.12. *Under the assumptions above, let $Z \subset M^\circ(E')$ be any irreducible component. Then the codimension of \overline{Z} in $\mathcal{H}_{d,g}^\dagger$ is at least $(s - r) + 1$.*

Proof. Let $z = \dim Z$. We use the associated scroll construction over Z . We have an open subset U of a vector bundle of rank $s + k + 1$ over Z and a morphism $U \rightarrow \mathcal{V}_g(\mathbf{F}_k, d\tau)$. Since $E' \neq E[k]$, the closure of the image of U is a proper subvariety of $\mathcal{V}_g(\mathbf{F}_k, \tau)$. In particular, we have $\dim U < \dim \mathcal{V}_g(\mathbf{F}_k, d\tau) = dk + 2d + g - 1$. The lemma follows from this inequality. \square

We now have the tools to determine all the Maroni divisors.

Proposition 2.13. *The Maroni locus $M(E) \subset \mathcal{H}_{d,g}$ is a divisor if and only if $g = (k - 1)(d - 1)$ for some integer $k \geq 1$, and $E = E[k - 1] = O(k - 1) \oplus O(k)^{\oplus d-3} \oplus O(k + 1)$. Furthermore, in this situation, $M(E[k - 1])$ is irreducible.*

Proof. If $\lfloor E \rfloor = k = \lfloor \frac{g+d-1}{d-1} \rfloor$, then the statement follows by applying Lemma 2.12. If, on the other hand, $\lfloor E \rfloor < \lfloor \frac{g+d-1}{d-1} \rfloor$, then the statement follows from statement 4 of Theorem 2.10. \square

We record a particularly interesting case of the irreducibility of the Maroni divisor.

Corollary 2.14. *Let $g = 2(d - 1)$. Then $M(E[2]) \subset \mathcal{H}_{d,g}$ is irreducible, and it is the ramification locus of the generically finite and dominant forgetful map $\mu: \mathcal{H}_{d,g} \rightarrow \mathcal{M}_g$.*

Proof. The irreducibility statement follows from Theorem 2.10. To show that $M(E[2])$ is the ramification locus of μ , consider $[\alpha: C \rightarrow \mathbf{P}^1] \in \mathcal{H}_{d,g}$ and the map of sheaves:

$$0 \rightarrow \alpha^*(T_{\mathbf{P}^1}) \rightarrow T_C \rightarrow N_\alpha \rightarrow 0.$$

The tangent space to $\mathcal{H}_{d,g}$ at α is $H^0(C, N_\alpha)/\alpha^*H^0(\mathbf{P}^1, T_{\mathbf{P}^1})$ and the tangent space to \mathcal{M}_g at C is $H^1(C, T_C)$. The map

$$d\mu: H^0(C, N_\alpha)/\alpha^*H^0(\mathbf{P}^1, T_{\mathbf{P}^1}) \rightarrow H^1(C, T_C)$$

fails to be surjective precisely when $H^1(C, \alpha^*T_{\mathbf{P}^1}) \neq 0$, that is, when $\alpha \in M(E[2])$. \square

2.2 Linear independence of T , D , and Δ

In this section, we prove that the divisorial components of the boundary of $\tilde{\mathcal{H}}_{d,g}$ are linearly independent. Define the closed loci T , D , Δ in $\tilde{\mathcal{H}}_{d,g}$ by

$$\begin{aligned} T &= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid \alpha^{-1}(q) = 3p_1 + p_2 + \cdots + p_{d-2} \text{ for some } q \text{ and distinct } p_i.\}} \\ D &= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid \alpha^{-1}(q) = 2p_1 + 2p_2 + p_3 + \cdots + p_{d-2} \text{ for some } q \text{ and distinct } p_i.\}} \\ \Delta &= \overline{\{[\alpha: C \rightarrow \mathbf{P}^1] \mid C \text{ is singular.}\}} \end{aligned}$$

These three loci correspond to the three possibilities of the limit when two branch points of a branched cover come together. Note that T , D , and Δ are irreducible and their union is the complement of $\tilde{\mathcal{H}}_{d,g}$ in $\mathcal{H}_{d,g}$.

Proposition 2.15. *For $d \geq 4$, the classes of T , D , and Δ are linearly independent in $\text{Pic}_{\mathbf{Q}}(\tilde{\mathcal{H}}_{d,g})$. For $d \geq 3$, the same is true for the classes of T and Δ .*

Proof. We construct curves with non-singular intersection matrix with our divisors. For this, a slight enlargement of $\tilde{\mathcal{H}}_{d,g}$ is more convenient. Define $\tilde{\mathcal{H}}_{d,g}^{ns}$ as the moduli space of $[\alpha: C \rightarrow \mathbf{P}^1]$ where C is an at worst nodal curve of arithmetic genus g , not necessarily irreducible, but without any separating nodes, and α is a map of degree d . The target \mathbf{P}^1 is taken to be unframed. It is easy to see that $\tilde{\mathcal{H}}_{d,g}$ is a dense open subset of $\tilde{\mathcal{H}}_{d,g}^{ns}$ with codimension two complement. Abusing notation, we denote the closures of T , D , and Δ in $\tilde{\mathcal{H}}_{d,g}^{ns}$ by the same letters. It suffices to prove the proposition for $\tilde{\mathcal{H}}_{d,g}^{ns}$.

We now construct test curves in $\tilde{\mathcal{H}}_{d,g}^{ns}$. Pick non-negative integers g_1 and g_2 with $g_1 + g_2 = g - 1$ and positive integers d_1 and d_2 with $d_1 + d_2 = d$. Take a family $\alpha_b: X_b \rightarrow \mathbf{P}^1$ of covers of degree d_1 and genus g_1 , where b denotes a parameter on a smooth complete curve B . Assume that we have two sections $p, q: B \rightarrow X$ with $\alpha_b(p_b) = 0$ and $\alpha_b(q_b) = \infty$ for all $b \in B$. Take $\beta: E \rightarrow \mathbf{P}^1$ to be a fixed simply branched cover of degree d_2 and genus g_2 , unramified over 0 and ∞ , and let $p', q' \in E$ be two points over 0 and ∞ respectively. Our test curve in $\tilde{\mathcal{H}}_{d,g}^{ns}$ is given by the family $\gamma_b: C_b \rightarrow \mathbf{P}^1$, where C_b is obtained by gluing (X_b, p_b, q_b) to the constant family (E, p', q') , and $\gamma_b: C_b \rightarrow \mathbf{P}^1$ is induced from $\alpha: X_b \rightarrow \mathbf{P}^1$ and $\beta: E \rightarrow \mathbf{P}^1$. The construction is depicted in Figure 1.




Figure 1: We construct families of covers parametrized by $b \in B$ by attaching a variable family of covers $\alpha_b: X_b \rightarrow \mathbf{P}^1$ to a fixed cover $\beta: E \rightarrow \mathbf{P}^1$.

Let T_α , D_α , and Δ_α denote the pullbacks of the divisor classes T , D , and Δ along the map from B to $\tilde{\mathcal{H}}_{d_1, g_1}$ given by α_b . Define T_γ , D_γ , and Δ_γ likewise. Let e be the intersection number of $\text{Br}(\alpha)$ with a horizontal section of $\mathbf{P}^1 \times B$. Denote by $[p]$ (resp. $[q]$) the class of $p(B)$ (resp. $q(B)$) on X .

Claim. With the notation above, we have

$$\deg T_\gamma = \deg T_\alpha + 3([p] + [q]) \cdot \text{Ram}(\alpha),$$

$$\deg D_\gamma = \deg D_\alpha + (2g_2 + 2d_2 - 2)e + 4e - 4([p] + [q]) \cdot \text{Ram}(\alpha), \text{ and}$$

$$\deg \Delta_\gamma = \deg \Delta_\alpha + [p]^2 + [q]^2.$$

Proof of the claim. The pullback of the line bundle $O(\Delta)$ from $\tilde{\mathcal{H}}_{d,g}^{ns}$ to B is given by

$$(N_{p/X} \otimes N_{p'/E}) \otimes (N_{q/E} \otimes N_{q'/E}) \otimes O_B(\Delta_\alpha),$$

where $N_{p/X}$ denotes the normal bundle of p in X , and so on. The third equation follows.

For a generic $b \in B$, the point of $\tilde{\mathcal{H}}_{d,g}^{ns}$ given by $\gamma_b: C_b \rightarrow \mathbf{P}^1$ does not lie in T or D . We have the following specializations:

1. $\alpha_b: X_b \rightarrow \mathbf{P}^1$ has a fiber of the form $3p_1 + p_2 + \dots$. Such b 's are precisely the points of T_α , each contributing 1 to $\deg T_\gamma$.
2. $\alpha_b: X_b \rightarrow \mathbf{P}^1$ has a fiber of the form $2p_1 + 2p_2 + p_3 + \dots$. Such b 's are precisely the points of D_α , each contributing 1 to $\deg D_\gamma$.
3. A branch point of $\alpha_b: X_b \rightarrow \mathbf{P}^1$ coincides with a branch point of $\beta: E \rightarrow \mathbf{P}^1$. There are $(2g_2 + 2d_2 - 2)e$ such b 's, each contributing 1 to $\deg D_\gamma$.
4. p_b (resp. q_b) is a ramification point of α_b . We compute the intersection multiplicity of B with T and D at such a point by looking at a versal deformation space of γ_b . We may restrict γ_b over an analytic neighborhood U of 0 (resp. ∞). Let x be a coordinate on U . Then $\gamma_b^{-1}(U) \rightarrow U$ has the form

$$U[y]/(y^3 - xy) \sqcup U \sqcup \dots \sqcup U \rightarrow U.$$

A versal deformation of this cover is given over $\text{Spec } \mathbf{C}[s, t]$ by

$$U[y]/(y^3 - xy - sx - t) \sqcup U \sqcup \dots \sqcup U \rightarrow U.$$

In $\text{Spec } \mathbf{C}[s, t]$, the divisor D does not contain the origin, and hence the intersection number of B with D at b is 0. The divisor $T \subset \text{Spec } \mathbf{C}[s, t]$ is defined by $t = 0$. The curve B approaches the origin along the locus where $U[y]/(y^3 - xy - sx - t)$ is singular, namely along $s^3 + t = 0$. We deduce that the intersection number of B with T at b is 3. There are $[p] \cdot \text{Ram}(\alpha)$ (resp. $[q] \cdot \text{Ram}(\alpha)$) such b 's, each contributing 3 to $\deg T_\gamma$.

5. p_b (resp. q_b) is not a ramification point of α_b , but lies over a branch point. Again, we look at a versal deformation of γ_b . In this case, $\gamma_b^{-1}(U) \rightarrow U$ has the form

$$U[y]/(y^2 - x) \sqcup U[z]/(z^2 - x^2) \sqcup U \sqcup \dots \sqcup U \rightarrow U.$$

A versal deformation of this cover is given over $\text{Spec } \mathbf{C}[s, t]$ by

$$U[y]/(y^2 - x) \sqcup U[z]/(z^2 - x^2 - sx - t) \sqcup U \sqcup \dots \sqcup U \rightarrow U.$$

In $\text{Spec } \mathbf{C}[s, t]$, the divisor T does not contain the origin, and hence the intersection number of B with T at b is 0. The divisor $D \subset \text{Spec } \mathbf{C}[s, t]$ is defined by $t = 0$. The curve B approaches the origin along the locus where $U[z]/(z^2 - x^2 - sx - t)$ is singular, namely along $s^2 - 4t = 0$. We deduce that the intersection number of B with D at b is 2. Let us count the number of such points, first for p_b , and analogously for q_b . The points b for which p_b is not a ramification point but lies over a branch point correspond to the intersection points of $\text{Br}(\alpha) \cap \{0\} \times B$ which are not the images of the points of $\text{Ram}(\alpha) \cap p(B)$. Note, however, that the image of a point of $\text{Ram}(\alpha) \cap p(B)$ is actually a point of tangency of $\text{Br}(\alpha)$ with $\{0\} \times B$, and hence contributes 2 to the intersection number $e = \text{Br}(\alpha) \cdot \{0\} \times B$. The remaining count, which we want, is therefore $e - 2[p] \cdot \text{Ram}(\alpha)$. Similarly, the count for q_b is $e - 2[q] \cdot \text{Ram}(\alpha)$.

The expressions for T_γ and D_γ follow from combining the above contributions. \square

Returning to the proof of the proposition, consider the following three particular test curves for $d \geq 4$.

B_1 : Take $\alpha_b: X_b \rightarrow \mathbf{P}^1$ to be a family of hyperelliptic curves of genus $g - 1$ obtained by taking a double cover $X \rightarrow \mathbf{P}^1 \times \mathbf{P}^1$ branched along a curve of type $(2g, 2)$. To have sections p and q of X over $\{0\} \times \mathbf{P}^1$ and $\{\infty\} \times \mathbf{P}^1$, let the branch divisor be tangent to $\{0\} \times \mathbf{P}^1$ and $\{\infty\} \times \mathbf{P}^1$. Take E to be a smooth rational curve and $\gamma: E \rightarrow \mathbf{P}^1$ a generic cover of degree $d - 2$.

B_2 : Take $\alpha_b: X_b \rightarrow \mathbf{P}^1$ to be a family of trigonal curves of genus $g - 1$ obtained by taking a general pencil on \mathbf{F}_0 in the linear system $|((g + 1)/2, 3)|$ if g is odd, or on \mathbf{F}_1 in the linear system $|3 \cdot \text{Directrix} + (g/2 + 2) \cdot \text{Fiber}|$ if g is even. Two base-points give p_b and q_b . Take E to be a rational curve and $\gamma: E \rightarrow \mathbf{P}^1$ a general cover of degree $d - 3$.

B_3 : Take $\alpha_b: X_b \rightarrow \mathbf{P}^1$ to be a family of hyperelliptic curves of genus $g - 2$ as in B_1 . Take E to be a smooth genus 1 curve and $\gamma: E \rightarrow \mathbf{P}^1$ a generic cover of degree $d - 2$. This curve exists only for $d \geq 4$.

Using the claim, we get the following non-singular intersection matrix.

	T	D	Δ
B_1	6	$4d - 12$	$8g - 6$
B_2	$3g + 9$	$8d - 24$	$7g - 3$
B_3	6	$4d - 8$	$8g - 14$

For $d = 3$, we take a pencil in \mathbf{F}_0 or \mathbf{F}_1 as in B_1 , but of trigonal curves of genus g , without any E . Then the middle column vanishes, and the second row becomes $(3g + 6, 0, 7g + 6)$, which is linearly independent from the first row. \square

3 Degree three

Let C be a curve of genus g and $\alpha: C \rightarrow \mathbf{P}^1$ a map of degree three. The relative canonical map embeds C as a divisor in a \mathbf{P}^1 -bundle $\mathbf{P}E$ over \mathbf{P}^1 , where E is a vector bundle of rank two and degree $g + 2$.

Let

$$E^{\text{gen}} = O\left(\left[\frac{g+2}{2}\right]\right) \oplus O\left(\left[\frac{g+2}{2}\right]\right)$$

be the most generic vector bundle on \mathbf{P}^1 of rank 2 and degree $g + 2$. Set

$$U_{E^{\text{gen}}} := \{\alpha \in \tilde{\mathcal{H}}_{3,g} \mid E_\alpha \cong E^{\text{gen}}\}.$$

Note that $U_{E^{\text{gen}}}$ is an open subset of $\tilde{\mathcal{H}}_{3,g}$.

Proposition 3.1. *The complement of $U_{E^{\text{gen}}}$ in $\tilde{\mathcal{H}}_{3,g}$ is a divisor if and only if g is even, in which case it is irreducible.*

Proof. This is the degree 3 case of Proposition 2.13. □

Let $\pi: \mathbf{P}E^{\text{gen}} \rightarrow \mathbf{P}^1$ be the projection. Set

$$V = H^0(\mathbf{P}^1, \text{Sym}^3 E^{\text{gen}} \otimes \det E^{\text{gen}\vee}).$$

Elements of $\mathbf{P}_{\text{sub}}V$ correspond to divisors in the linear series of the line bundle $O_{\mathbf{P}E^{\text{gen}}}(3) \otimes \pi^*(\det E^{\text{gen}})^\vee$ on $\mathbf{P}E^{\text{gen}}$. Let $C_v \subset \mathbf{P}E^{\text{gen}}$ be the divisor corresponding to $v \in V$. Let $V^\circ \subset \mathbf{P}_{\text{sub}}V$ be the open locus consisting of $v \in V^\circ$ for which C_v is irreducible and at worst nodal. Let $G := \text{Aut}(\pi)$ be the group of automorphisms of $\mathbf{P}E^{\text{gen}}$ over \mathbf{P}^1 . Then G acts on V° . The assignment

$$v \mapsto [\pi: C_v \rightarrow \mathbf{P}^1]$$

gives a map

$$q: V^\circ \rightarrow \tilde{\mathcal{H}}_{3,g}^\dagger.$$

Denote by $U_{E^{\text{gen}}}^\dagger$ the preimage of $U_{E^{\text{gen}}}$ under $\tilde{\mathcal{H}}_{3,g}^\dagger \rightarrow \tilde{\mathcal{H}}_{3,g}$.

Proposition 3.2. *The image of q is $U_{E^{\text{gen}}}^\dagger$. The fibers of q consist of single G -orbits.*

Proof. For brevity, set $E = E^{\text{gen}}$. For $v \in V^\circ$, consider the sequence

$$0 \rightarrow O_{\mathbf{P}E}(-3) \otimes \pi^* \det E \rightarrow O_{\mathbf{P}E} \rightarrow O_{C_v} \rightarrow 0.$$

Applying $R\pi_*$, we get

$$0 \rightarrow O_{\mathbf{P}^1} \rightarrow \pi_* O_{C_u} \rightarrow E^\vee \rightarrow 0, \tag{3.1}$$

which says that the Tschirnhausen bundle of $C_u \rightarrow \mathbf{P}^1$ is E . Conversely, from the Casnati–Ekedahl resolution, it follows that every point of $U_{E^{\text{gen}}}^{\dagger}$ is in the image of q .

Let $u, v \in U_{E^{\text{gen}}}^{\dagger}$ be in a fiber of q . Then there is an isomorphism $C_u \rightarrow C_v$ over the identity of \mathbf{P}^1 . The sequence (3.1) for C_u and C_v shows that such an isomorphism induces an isomorphism $E \rightarrow E$. The induced automorphism of $\mathbf{P}E$ over \mathbf{P}^1 takes C_u to C_v and hence u to v . \square

Proposition 3.3. *[Picard rank conjecture for degree three] We have $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{3,g} = 0$.*

Proof. Retain the notation introduced above. For brevity, set $U = U_{E^{\text{gen}}}$ and $U^{\dagger} = U_{E^{\text{gen}}}^{\dagger}$. Then $V^{\circ} \rightarrow U^{\dagger}$ is a quotient by G and $U^{\dagger} \rightarrow U$ is a quotient by PGL_2 . By Proposition 1.2 and Proposition 3.2, we have

$$\begin{aligned} \text{rk Pic}_{\mathbf{Q}} U &\leq \text{rk Pic}_{\mathbf{Q}} U^{\dagger} + \text{rk } \chi(\text{PGL}_2) = \text{rk Pic}_{\mathbf{Q}} U^{\dagger} \\ &\leq \text{rk Pic}_{\mathbf{Q}} V^{\circ} + \text{rk } \chi(G) \leq 1 + \text{rk } \chi(G). \end{aligned}$$

The final inequality follows because V° is an open subset of a projective space. Let e be the number of divisorial components of $\tilde{\mathcal{H}}_{3,g} \setminus U$. We then get the bound

$$\text{rk Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{3,g} \leq \text{rk Pic}_{\mathbf{Q}} U + e \leq 1 + \text{rk } \chi(G) + e.$$

If g is even, then

$$\begin{aligned} G &= \text{PGL}_2 \\ \text{rk } \chi(G) &= 0 \\ e &= 1 \quad \text{by Proposition 3.1.} \end{aligned}$$

If g is odd, then

$$\begin{aligned} G &= \left\{ \begin{pmatrix} a & l \\ & b \end{pmatrix} \mid a, b \in \mathbf{C}^*, l \in H^0(\mathbf{P}^1, \mathcal{O}(1)) \right\} / \mathbf{C}^* \\ \text{rk } \chi(G) &= 1 \\ e &= 0 \quad \text{by Proposition 3.1.} \end{aligned}$$

In either case, we have

$$\text{rk Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{3,g} \leq 2.$$

By Proposition 2.15, the classes in $\text{Pic}_{\mathbf{Q}}(\tilde{\mathcal{H}}_{3,g})$ of the two components of $\tilde{\mathcal{H}}_{3,g} \setminus \mathcal{H}_{3,g}$ are linearly independent. Therefore, we get $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{3,g} = 0$ as desired. \square

4 Degree four

Let C be a curve of genus g and $\alpha: C \rightarrow \mathbf{P}^1$ a map of degree four. The relative canonical map embeds C into a \mathbf{P}^2 -bundle $\mathbf{P}E$ over \mathbf{P}^1 , where E is a vector bundle of rank three and degree $g+3$. The Casnati-Ekedahl structure theorem provides the following resolution of O_C :

$$0 \rightarrow \pi^* \det E(-4) \rightarrow \pi^* F(-2) \rightarrow O_{\mathbf{P}E} \rightarrow O_C \rightarrow 0,$$

where F is a vector bundle of rank two and degree $g+3$.

Explicitly, we can describe $C \subset \mathbf{P}E$ as follows. Write $F = O(a) \oplus O(b)$, where $a+b = g+3$, and $a \leq b$. Let h denote the divisor class associated to $O_{\mathbf{P}E}(1)$ on $\mathbf{P}E$ and f the class of the fiber of the projection $\pi: \mathbf{P}E \rightarrow \mathbf{P}^1$. Then the curve C is the complete intersection of two divisors

$$C = Q_a \cap Q_b,$$

where $[Q_a] = 2h - af$ and $[Q_b] = 2h - bf$.

Even more explicitly, we can describe the equations of Q_a and Q_b as follows. Write $E = O(m_1) \oplus O(m_2) \oplus O(m_3)$. Over an open set $U \subset \mathbf{P}^1$, let X, Y , and Z denote the relative coordinates on $\mathbf{P}E|_U$ corresponding to the three summands of E . Assume that $m_1 \leq m_2 \leq m_3$. Over U , the divisor Q_a is the zero locus of a form

$$p_{1,1}X^2 + p_{1,2}XY + p_{1,3}XZ + p_{2,2}Y^2 + p_{2,3}YZ + p_{3,3}Z^2 \quad (4.1)$$

where $p_{i,j}$ is the restriction to U of a global section of $O(m_i + m_j - a)$. Similarly, over U , the divisor Q_b is the zero locus of a form

$$q_{1,1}X^2 + q_{1,2}XY + q_{1,3}XZ + q_{2,2}Y^2 + q_{2,3}YZ + q_{3,3}Z^2 \quad (4.2)$$

where $q_{i,j}$ is the restriction to U of a global section of $O(m_i + m_j - b)$.

The irreducibility of C puts some restrictions on the possible (E, F) . Indeed, if $p_{1,1} = q_{1,1} = 0$, then the section $[X : Y : Z] = [1 : 0 : 0]$ of $\mathbf{P}E$ is contained in both Q_a and Q_b , making $C = Q_a \cap Q_b$ reducible. An irreducible C thus forces

$$2m_1 \geq a. \quad (4.3)$$

Proposition 4.1. *Let E be a vector bundle of rank 3 and degree $g+3$ and F a vector bundle of rank 2 and degree $g+3$. If the locus $M(E, F)$ is non-empty, then it is irreducible and unirational.*

Proof. Consider the dense open subset $M^\circ(E, F) \subset M(E, F)$ corresponding to $\alpha \in \mathcal{H}_{4,g}$ that have $E_\alpha \cong E$ and $F_\alpha \cong F$. It suffices to prove the statement for $M^\circ(E, F)$.

Consider the vector space

$$V := H^0(\mathbf{P}^1, F^\vee \otimes \text{Sym}^2 E).$$

Elements of V correspond to maps $\pi^*F(-2) \rightarrow \mathcal{O}_{\mathbf{P}^E}$. Let $V^\circ \subset V$ be the open subset where the ideal generated by the image of $\pi^*F(-2)$ defines a smooth curve, simply branched over \mathbf{P}^1 . Then V° surjects onto $M^\circ(E, F)$. \square

Remark 4.2. From the dominant map $V^\circ \rightarrow M(E, F)$ in the proof of Proposition 4.1, it is easy to compute the codimension of $M(E, F)$ in $\mathcal{H}_{4,g}$, which is

$$\text{codim } M(E, F) = \dim \text{Ext}^1(E, E) + \dim \text{Ext}^1(F, F) - \dim \text{Ext}^1(F, \text{Sym}^2 F).$$

We may think of $\dim \text{Ext}^1(E, E) + \dim \text{Ext}^1(F, F)$ as the ‘expected codimension.’ The next example shows that the actual codimension is not always the expected codimension.

Example 4.3. Let $E = O(m) \oplus O(2m) \oplus O(g + 3 - 3m)$, where $\lceil \frac{g+3}{6} \rceil \leq m < \frac{g+3}{5}$. To get an irreducible curve C , the only possibility for F is $F = O(2m) \oplus O(g + 3 - 2m)$, by (4.3). The resulting locus $M(E, F)$ is not of expected codimension because $\dim \text{Ext}^1(F, \text{Sym}^2 E)$ is nonzero.

Example 4.4. The Maroni locus $M(E)$ may be reducible. Let $g = 12$, and consider the bundle $E = O(3) \oplus O(5) \oplus O(7)$. Then the reader can easily check (using Bertini’s theorem) that $M(E, F)$ and $M(E, F')$ are nonempty and of *equal* codimension $\dim \text{Ext}^1(E, E)$ for the bundles $F = O(6) \oplus O(9)$ and $F' = O(5) \oplus O(10)$. Therefore $M(E, F)$ and $M(E, F')$ are two components of $M(E)$. It is easy to see by analyzing the explicit equations that these are the only components of $M(E)$.

Let E^{gen} (resp. F^{gen}) be the most generic vector bundle on \mathbf{P}^1 of rank 3 (resp. 2) and degree $g + 3$. Define

$$\begin{aligned} U_{E^{\text{gen}}} &:= \{\alpha \in \tilde{\mathcal{H}}_{4,g} \mid E_\alpha \cong E^{\text{gen}}\}, \\ U_{F^{\text{gen}}} &:= \{\alpha \in \tilde{\mathcal{H}}_{4,g} \mid F_\alpha \cong F^{\text{gen}}\}, \\ U_{E^{\text{gen}}, F^{\text{gen}}} &:= U_{E^{\text{gen}}} \cap U_{F^{\text{gen}}}. \end{aligned}$$

It is easy to see that these are open subsets of $\tilde{\mathcal{H}}_{d,g}$. Our next task is to identify the divisorial components of their complements.

Proposition 4.5. *The subvariety $M := \tilde{\mathcal{H}}_{4,g} \setminus U_{E^{\text{gen}}}$ is a divisor if and only if g is divisible by three, in which case it is irreducible.*

Proof. This is the degree 4 case of Proposition 2.13. \square

For the complement of $U_{F^{\text{gen}}}$, we could do a careful analysis of the defining equations of C in \mathbf{P}^E , as we will have to do for the next case of $d = 5$. But we can take a more geometric approach using the *resolvent cubic construction*. Originally due to Recillas [Rec73], the construction can be described as follows. For simplicity, we give an informal

description, restricting to simply branched covers. See [Cas98] for a detailed account. Consider a point $[\alpha: C \rightarrow \mathbf{P}^1]$ of $\mathcal{H}_{4,g}$. The resolution of O_C as an $O_{\mathbf{P}E_\alpha}$ module shows that $C \subset \mathbf{P}E_\alpha$ is the complete intersection of two relative quadrics. A fiber of $\mathbf{P}F_\alpha \rightarrow \mathbf{P}^1$ naturally corresponds to the pencil of conics in the corresponding fiber of $\mathbf{P}E_\alpha \rightarrow \mathbf{P}^1$ containing the corresponding fiber of $C \rightarrow \mathbf{P}^1$. Each such pencil contains three singular conics, counted with multiplicity. The total locus of these singular conics forms a trigonal curve $R(C) \subset \mathbf{P}F_\alpha$. Let $R(\alpha): R(C) \rightarrow \mathbf{P}^1$ be the projection. We call $R(\alpha)$ the *resolvent cubic* of α . Using that $C \rightarrow \mathbf{P}^1$ is simply branched, it is easy to check that $R(C)$ is smooth and the branch divisor of $R(\alpha)$ coincides with the branch divisor of α . In particular, $R(C)$ has genus $g + 1$. The association $\alpha \rightarrow R(\alpha)$ defines a map

$$R: \mathcal{H}_{4,g} \rightarrow \mathcal{H}_{3,g+1},$$

which we call the *resolvent cubic map*. The fiber of R over a point $[D \rightarrow \mathbf{P}^1] \in \mathcal{H}_{3,g+1}$ corresponds bijectively to the set of étale double covers $D' \rightarrow D$ (see [Rec73] or [Cas98, Theorem 6.5]). In particular, R is a finite morphism.

Proposition 4.6. *Let F be a vector bundle of rank 2 and degree $g + 3$ on \mathbf{P}^1 . The Casnati-Ekedahl locus $C(F) \subset \mathcal{H}_{4,g}$ is non-empty if and only if $[F] \geq \lceil \frac{g+3}{3} \rceil$. In this case, it is of the expected codimension $\dim \text{Ext}^1(F, F)$.*

Proof. Consider a point $[\alpha: C \rightarrow \mathbf{P}^1]$ of $\mathcal{H}_{4,g}$ and its resolvent cubic $R(\alpha): R(C) \rightarrow \mathbf{P}^1$. Since $R(C) \subset \mathbf{P}F_\alpha$, and F_α is a vector bundle of rank two and degree $(g + 1) + 2$, it must be the Tschirnhausen bundle of $R(C)$. That is, we have $E_{R(\alpha)} = F_\alpha$. By [Rec73], the map R is finite, and hence $C(F) = R^{-1}(M(F))$. Both of the statements about $C(F)$ now follow from the corresponding statements about $M(F)$. \square

Proposition 4.7. *Let $g \geq 4$. The subvariety $CE := \mathcal{H}_{4,g} \setminus U_{F^{\text{gen}}}$ has codimension at least two if g is even and is an irreducible divisor if g is odd.*

Proof. The image $R(U_{F^{\text{gen}}}) \subset \mathcal{H}_{3,g+1}$ is the open locus of trigonal covers having F^{gen} as their Tschirnhausen bundle. The complement $Z := \mathcal{H}_{3,g+1} \setminus R(U_{F^{\text{gen}}})$ has codimension at least two if $g + 1$ is odd and it is the Maroni divisor if $g + 1$ is even (Proposition 3.1). The complement $\mathcal{H}_{4,g} \setminus U_{F^{\text{gen}}}$ is the preimage $R^{-1}(Z)$. Therefore, the statements about the codimension follow from the finiteness of R .

For the question of reducibility, let $F = O(k - 1) \oplus O(k + 1)$ with $k = (g + 3)/2 \geq 3$. The claim is that $C(F)$ is irreducible when $g > 3$, and has two components when $g = 3$. We have

$$C(F) = \bigcup_E M(E, F).$$

By Proposition 4.1, the varieties $M(E, F)$ are irreducible. Therefore, every component of $C(F)$ must be of the form $M(E, F)$ for some E .

Let $g > 3$ and suppose $E \neq E^{\text{gen}}$. The inclusion $M(E, F) \subset M(E)$ and Proposition 2.13 imply that $M(E, F)$ is a divisor if and only if $M(E, F) = M(E)$ and $E = O(m-1) \oplus O(m) \oplus O(m+1)$. By choosing two generic quadrics as in (4.1) and (4.2), we can explicitly construct a curve in $M(E, F^{\text{gen}})$, showing that $M(E, F) \neq M(E)$. Thus, it follows that the only component of $C(F)$ is $M(E^{\text{gen}}, F)$. \square

Example 4.8. The divisor $\mathcal{H}_{4,g} \setminus U_{F^{\text{gen}}}$ is not irreducible for $g = 3$. Indeed, take $F = O(2) \oplus O(4)$. Then $M(E^{\text{gen}}, F)$ is an irreducible component. Now consider the only other possibility for E , namely $E = O(1) \oplus O(2) \oplus O(3)$. By (4.3), a cover in $M(E)$ must have $F = O(2) \oplus O(4)$. Furthermore, for this E and F , we can choose the two quadrics generically and see that $M(E, F)$ is nonempty. Therefore, $M(E) = M(E, F)$ is another component of $\mathcal{H}_{4,g} \setminus U_{F^{\text{gen}}}$.

Our next goal is to exhibit $U_{E^{\text{gen}}, F^{\text{gen}}}$ as a quotient. Let $\pi: \mathbf{P}E^{\text{gen}} \rightarrow \mathbf{P}^1$ be the projection. For brevity, set $E = E^{\text{gen}}$ and $F = F^{\text{gen}}$. Set

$$V := H^0(\mathbf{P}^1, F^\vee \otimes \text{Sym}^2 E).$$

An element $v \in \mathbf{P}_{\text{sub}}V$ corresponds to a map $\pi^*F(-2) \rightarrow O_{\mathbf{P}E}$. Let C_v be the zero locus of the image of this map. Let $V^\circ \subset \mathbf{P}_{\text{sub}}V$ be the open locus consisting of $v \in \mathbf{P}_{\text{sub}}V$ for which C_v is irreducible and at worst nodal. Let $G_F := \text{Aut}(\mathbf{P}F/\mathbf{P}^1)$ and $G_E := \text{Aut}(\mathbf{P}E/\mathbf{P}^1)$. Then $G_F \times G_E$ acts on V° . The assignment

$$v \mapsto [\pi: C_v \rightarrow \mathbf{P}^1]$$

defines a map

$$q: V^\circ \rightarrow \tilde{\mathcal{H}}_{4,g}^\dagger.$$

Denote by $U_{E,F}^\dagger$ the preimage of $U_{E,F}$ under $\tilde{\mathcal{H}}_{4,g}^\dagger \rightarrow \tilde{\mathcal{H}}_{4,g}$.

Proposition 4.9. *The image of q is $U_{E^{\text{gen}}, F^{\text{gen}}}^\dagger$. The fibers of q consist of single G -orbits.*

Proof. The proof is exactly analogous to the proof of Proposition 3.2. \square

Proposition 4.10. *[Picard rank conjecture for degree four] We have $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{4,g} = 0$.*

Proof. Retain the notation introduced above. For brevity, set $U = U_{E^{\text{gen}}, F^{\text{gen}}}$ and $U^\dagger = U_{E^{\text{gen}}, F^{\text{gen}}}^\dagger$. By Proposition 1.2 and Proposition 4.9, we have

$$\begin{aligned} \text{rk Pic}_{\mathbf{Q}} U &\leq \text{rk Pic}_{\mathbf{Q}} U^\dagger + \text{rk } \chi(\text{PGL}_2) = \text{rk Pic}_{\mathbf{Q}} U^\dagger \\ &\leq \text{rk Pic}_{\mathbf{Q}} V^\circ + \text{rk } \chi(G) \leq 1 + \text{rk } \chi(G). \end{aligned}$$

The final inequality follows because V° is an open subset of a projective space. Let e be the number of divisorial components of $\tilde{\mathcal{H}}_{4,g} \setminus U$. We then get the bound

$$\text{rk Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{4,g} \leq \text{rk Pic}_{\mathbf{Q}} U + e \leq 1 + \text{rk } \chi(G) + e.$$

Recall that $G = G_{F^{\text{gen}}} \times G_{E^{\text{gen}}}$.

If g is an odd multiple of 3, then

$$G = \text{PGL}_2 \times \text{PGL}_3$$

$$\text{rk } \chi(G) = 0$$

$e = 2$ corresponding to M in Proposition 4.5 and CE in Proposition 4.7.

If g is odd, but not divisible by 3, then

$$G = \text{PGL}_2 \times G_E$$

$$G_E = \left\{ \begin{pmatrix} a & b & l_1 \\ c & d & l_2 \\ 0 & 0 & e \end{pmatrix} \mid a, b, c, d, e \in \mathbf{C}, e(ad - bc) \in \mathbf{C}^*, l_i \in H^0(\mathbf{P}^1, \mathcal{O}(1)) \right\} / \mathbf{C}^*.$$

$$\text{rk } \chi(G) = 1$$

$e = 1$ corresponding to CE in Proposition 4.7.

If g is even and divisible by 3, then

$$G = G_F \times \text{PGL}_2$$

$$G_F = \left\{ \begin{pmatrix} a & l \\ & b \end{pmatrix} \mid a, b \in \mathbf{C}^*, l \in H^0(\mathbf{P}^1, \mathcal{O}(1)) \right\} / \mathbf{C}^*$$

$$\text{rk } \chi(G) = 1$$

$e = 1$ corresponding to M in Proposition 4.5.

If g is even and not divisible by 3, then

$G = G_F \times G_E$ where G_F and G_E are as in the previous two cases,

$$\text{rk } \chi(G) = 2$$

$$e = 0.$$

In all cases, we get

$$\text{rk Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{4,g} \leq 3.$$

By Proposition 2.15, the classes in $\text{Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{4,g}$ of the three components of $\tilde{\mathcal{H}}_{4,g} \setminus \mathcal{H}_{4,g}$ are linearly independent. Therefore, we get $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{4,g} = 0$ as desired. \square

5 Degree five

Let C be a curve of genus g and $\alpha: C \rightarrow \mathbf{P}^1$ a map of degree five. The relative canonical map embeds C into a \mathbf{P}^3 bundle $\mathbf{P}E$ over \mathbf{P}^1 , where E is a vector bundle of rank four and

degree $g + 4$. The Casnati-Ekedahl structure theorem provides the following resolution of O_C :

$$0 \rightarrow \pi^* \det E(-5) \rightarrow \pi^*(F^\vee(\det E))(-3) \rightarrow \pi^*F(-2) \rightarrow O_{\mathbf{P}E} \rightarrow O_C \rightarrow 0$$

where F is a vector bundle of rank three and degree $2g + 8$.

Explicitly, we can describe $C \subset \mathbf{P}E$ as follows. The resolution is determined completely by the middle map

$$w: \pi^*(F^\vee(\det E))(-3) \rightarrow \pi^*F(-2).$$

This map may be viewed an element of the vector space $H^0(\mathbf{P}^1, F \otimes F \otimes E(-\det E))$. Due to a theorem of Casnati [Cas96], w can be taken to be anti-symmetric, that is, in the subspace

$$V := H^0(\mathbf{P}^1, \wedge^2 F \otimes E \otimes \det E^\vee).$$

Even more explicitly, we can describe the defining equations of C as follows. Let

$$\begin{aligned} F &= O(n_1) \oplus \cdots \oplus O(n_5), \text{ where } n_1 \leq \cdots \leq n_5, \text{ and} \\ E &= O(m_1) \oplus \cdots \oplus O(m_4), \text{ where } m_1 \leq \cdots \leq m_4. \end{aligned}$$

We represent an element $w \in V$ by a skew symmetric matrix of forms

$$M_w = \begin{pmatrix} 0 & L_{1,2} & L_{1,3} & L_{1,4} & L_{1,5} \\ -L_{1,2} & 0 & L_{2,3} & L_{2,4} & L_{2,5} \\ -L_{1,3} & -L_{2,3} & 0 & L_{3,4} & L_{3,5} \\ -L_{1,4} & -L_{2,4} & -L_{3,4} & 0 & L_{4,5} \\ -L_{1,5} & -L_{2,5} & -L_{3,5} & -L_{4,5} & 0 \end{pmatrix} \quad (5.1)$$

where $L_{i,j} \in H^0(\mathbf{P}^1, E \otimes \det E^\vee \otimes O(n_i + n_j))$. In $\mathbf{P}E$, the curve C_w is cut out by the 4×4 sub-Pfaffians of the matrix M_w .

The irreducibility of C puts some restrictions on the possible matrices. Indeed, suppose

$$L_{1,2} = L_{1,3} = 0.$$

Then the Pfaffian Q_5 of the submatrix obtained by eliminating the fifth row and column is

$$Q_5 = L_{1,2}L_{3,4} - L_{1,3}L_{2,4} + L_{2,3}L_{1,4} = L_{2,3}L_{1,4}.$$

Since Q_5 is reducible, it forces C_w to be reducible.

Suppose further that $E = O(k)^r \oplus O(k+1)^{4-r}$, where $0 \leq r \leq 3$. Then the observation above implies that the maximum of the degrees of the summands of $E \otimes (\det E^\vee) \otimes O(n_1 + n_3)$ must be nonnegative, meaning

$$n_1 + n_3 + k - (g + 4) \geq -1.$$

Since the n_i are increasing, we get the inequalities

$$n_i + n_j + (k + 1) - (g + 4) \geq 0 \text{ for every } (i, j) \text{ with } i \neq j \text{ except } (i, j) = (1, 2). \quad (5.2)$$

Let E^{gen} (resp. F^{gen}) be the most generic vector bundle on \mathbf{P}^1 of rank 4 (resp. 5) and degree $g + 4$ (resp. $2g + 8$). Define $U_{E^{\text{gen}}}$, $U_{F^{\text{gen}}}$, and $U_{E^{\text{gen}}, F^{\text{gen}}}$ as before. These are the open subsets of $\tilde{\mathcal{H}}_{5,g}$ consisting of covers α for which E_α, F_α , and both E_α and F_α are the most generic.

Proposition 5.1. *The subvariety $M := \tilde{\mathcal{H}}_{5,g} \setminus U_{E^{\text{gen}}}$ has codimension at least two if g is not divisible by 4, and has a unique divisorial component if g is divisible by 4.*

Proof. This is the degree 5 case of Proposition 2.13. □

For the complement of $U_{F^{\text{gen}}}$, we must analyze the defining equations of C in $\mathbf{P}E$.

Proposition 5.2. *The subvariety $CE := \mathcal{H}_{5,g} \setminus U_{F^{\text{gen}}}$ has codimension at least two if $g + 4$ is not a multiple of 5 (with the exception of $g = 3$ in which case the complement parametrizes hyperelliptic curves), and contains a unique divisorial component if $g + 4$ is a multiple of 5.*

Proof. We must characterize the Casnati-Ekedahl loci $C(F)$ which are divisorial. We have

$$C(F) = \bigcup_E M(E, F).$$

The loci $M(E, F)$ are irreducible by virtually the same argument as in Proposition 4.1 (In the proof, just take $V = H^0(\mathbf{P}^1, \wedge^2 F \otimes E \otimes \det E^\vee)$.) Therefore, any component of $C(F)$ must be of the form $M(E, F)$. From the explicit description of degree 5 covers above, it is straightforward to compute that

$$\text{codim } M(E, F) = \dim \text{Ext}^1(E, E) + \dim \text{Ext}^1(F, F) - h^1(\wedge^2 F \otimes E \otimes \det E^\vee).$$

Suppose $E \neq E^{\text{gen}}$. Then $M(E, F) \subset M(E)$. By Proposition 2.13, $M(E)$ has codimension at least two unless $E = O(k) \oplus O(k+1)^{\oplus d-3} \oplus O(k+2)$. In this case, using the explicit description of degree 5 covers, it is easy to construct covers α with $E_\alpha = E$ and $F_\alpha = F^{\text{gen}}$. Thus, $M(E, F) \neq M(E)$, and since $M(E)$ is irreducible, $M(E, F) \subset M(E)$ has codimension at least one. Therefore, $M(E, F) \subset \mathcal{H}_{4,g}$ has codimension at least two.

Therefore, for $M(E, F)$ to be divisorial, we must have $E = E^{\text{gen}}$. In this case, we have

$$\text{codim } M(E, F) = \dim \text{Ext}^1(F, F) - h^1(\wedge^2 F \otimes E \otimes \det E^\vee).$$

Suppose $h^1(\wedge^2 F \otimes E \otimes \det E^\vee) = 0$. Note that $\dim \text{Ext}^1(F, F) = 1$ if and only if

$$F = O(n-1) \oplus O(n) \oplus O(n) \oplus O(n) \oplus O(n+1).$$

In this case $5n = 2(g + 4)$, and hence 5 divides $g + 4$.

We are thus reduced to showing that $M(E, F)$ is not a divisor when $E = E^{\text{gen}}$ and

$$h^1(\wedge^2 F \otimes E(-\det E)) > 0,$$

with the exception of $g = 3$. Write

$$E = O(k)^{\oplus r} \oplus O(k+1)^{\oplus 4-r} \text{ where } 0 \leq r \leq 3,$$

and

$$F = O(n_1) \oplus O(n_2) \oplus O(n_3) \oplus O(n_4) \oplus O(n_5), \text{ where } n_1 \leq \dots \leq n_5.$$

Consider an anti-symmetric matrix

$$M_w = (L_{i,j}) \quad 1 \leq i, j \leq 5,$$

as in (5.1), representing an element of $H^0(\wedge^2 F \otimes E \otimes \det E^\vee)$. Inequality (5.2) implies that any contribution to $h^1(\wedge^2 F \otimes E \otimes \det E^\vee)$ must come from the $L_{1,2}$ entry. In other words, we have

$$h^1(\wedge^2 F \otimes E(-\det E)) = h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)).$$

Since $E = E^{\text{gen}}$, we have $h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)) > 0$ if and only if

$$n_1 + n_2 + (k+1) - (g+4) < 0.$$

Hence, we get

$$\begin{aligned} h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)) &= 4(-(n_1 + n_2 + k - (g+4)) - 1) - (4-r) \\ &= 4g - 4(n_1 + n_2 + k) + r + 8. \end{aligned}$$

Equation (5.2) tells us that $n_1 + n_3 + (k+1) - (g+4) \geq 0$, which implies $n_2 < n_3$. Therefore,

$$\dim \text{Ext}^1(F, F) \geq (2n_5 + 2n_4 + 2n_3) - 3(n_1 + n_2) - 6.$$

Combining the two, we get

$$\dim \text{Ext}^1(F, F) - h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)) \geq 2n_5 + 2n_4 + 2n_3 + n_1 + n_2 - 3(g+4) - 2.$$

Using $n_1 + \dots + n_5 = 2(g+4)$, the above inequality becomes

$$\dim \text{Ext}^1(F, F) - h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)) \geq (g+4) - (n_1 + n_2) - 2.$$

Finally, by using the assumption $n_1 + n_2 + (k+1) - (g+4) < 0$, we conclude that

$$\text{codim } M(E^{\text{gen}}, F) = \dim \text{Ext}^1(F, F) - h^1(E \otimes \det E^\vee \otimes O(n_1 + n_2)) > k - 1.$$

If $k > 1$, then we get $\text{codim } M(E^{\text{gen}}, F) > 1$ as desired. We consider the cases where $k = 1$ on an individual basis. These cases correspond to $0 \leq g \leq 4$.

Case: $g = 4$. Then $E^{\text{gen}} = O(2)^{\oplus 4}$ and $F^{\text{gen}} = O(3)^{\oplus 4} \oplus O(4)$. The relative canonical map embeds C in $\mathbf{P}E^{\text{gen}} \simeq \mathbf{P}^3 \times \mathbf{P}^1$. The projection to \mathbf{P}^3 restricts to the canonical map on C . Therefore, if C is non-hyperelliptic, then there is only one quadric in \mathbf{P}^3 containing the canonical model of C . This means that the bundle F has exactly one $O(4)$ summand, and hence $F \cong F^{\text{gen}}$. The locus where C is hyperelliptic is easily seen to be codimension 2 in $\mathcal{H}_{5,4}$. This exhausts all possibilities in this case.

Case: $g = 3$. Then $E^{\text{gen}} = O(1) \oplus O(2)^{\oplus 3}$ and $F^{\text{gen}} = O(2) \oplus O(3)^{\oplus 4}$. Consider the special bundle $F = O(2) \oplus O(2) \oplus O(3)^{\oplus 2} \oplus O(4)$. Then

$$\dim \text{Ext}^1(F, F) - h^1(\wedge^2 F \otimes E \otimes \det E^\vee) = 1.$$

Now consider a general $[\alpha: C \rightarrow \mathbf{P}^1] \in M(E, F) \subset \mathcal{H}_{5,3}$. Let $[X : Y : Z : W]$ denote the homogeneous coordinates (locally over \mathbf{P}^1) on $\mathbf{P}E$ corresponding to the summands of E . As usual, denote by h the class of $O_{\mathbf{P}E}(1)$ and by f the class of the fiber of $\mathbf{P}E \rightarrow \mathbf{P}^1$. Since $O(4)$ is a summand of F , there exists a unique effective divisor Q of class $2h - 4f$ on $\mathbf{P}E$ which contains C . The quadric Q may be written as the zero locus of a form

$$c_0 Y^2 + c_1 YZ + \cdots + c_5 W^2,$$

where c_i are constants. Let $p: \mathbf{P}E \dashrightarrow \mathbf{P}^2 \times \mathbf{P}^1$ be the projection from the section $[1 : 0 : 0 : 0]$, and $g: \mathbf{P}E \dashrightarrow \mathbf{P}^2 \times \mathbf{P}^1 \rightarrow \mathbf{P}^2$ the composition with the projection onto the first factor. Then the rational map g is given by the linear system $|h - 2f|$ on $\mathbf{P}E$, which restricts to the canonical series on C . However, the fact that C lies on the relative quadric Q means that the image $g(C)$ is exactly the conic defined by the equation for Q . Thus, C is hyperelliptic. Given the above geometric understanding of the $O(4)$ summand of F , it is easy to show that if we begin with a hyperelliptic curve C , and a degree 5 map $\alpha: C \rightarrow \mathbf{P}^1$, then F_α must contain a unique $O(4)$ summand. By the inequalities in (5.2), there are no other choices for F .

Case: $g = 1, 2$. In these cases, we leave it to the reader to see that there are no nontrivial Casnati-Ekedahl or Maroni loci. □

As before, we now exhibit $U_{E^{\text{gen}}, F^{\text{gen}}}$ as a quotient. For brevity, set $E = E^{\text{gen}}$ and $F = F^{\text{gen}}$. Set

$$V := H^0(\mathbf{P}^1, \wedge^2 F \otimes E \otimes \det E).$$

An element $v \in \mathbf{P}_{\text{sub}}V$ defines an anti-symmetric matrix as in (5.1). Let C_v be the zero locus of the 4×4 sub-Pfaffians of this matrix. Let $V^\circ \subset \mathbf{P}_{\text{sub}}V$ be the open locus consisting of v for which C_v is irreducible and at worst nodal. Let $G_F := \text{Aut}(\mathbf{P}F/\mathbf{P}^1)$ and $G_E := \text{Aut}(\mathbf{P}E/\mathbf{P}^1)$. Then $G := G_F \times G_E$ acts on V° . The assignment $v \mapsto [\pi: C_v \rightarrow \mathbf{P}^1]$ defines a map

$$q: V^\circ \rightarrow \widetilde{\mathcal{H}}_{5,g}^\dagger.$$

Let $U_{E,F}^\dagger$ be the preimage of $U_{E^{\text{gen}},F^{\text{gen}}}$ under $\tilde{\mathcal{H}}_{5,g}^\dagger \rightarrow \tilde{\mathcal{H}}_{5,g}$.

Proposition 5.3. *The image of q is $U_{E^{\text{gen}},F^{\text{gen}}}^\dagger$. The fibers of q consist of single G -orbits.*

Proof. The proof is exactly analogous to that of Proposition 3.2. □

Proposition 5.4. *[Picard rank conjecture for degree five] We have $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{5,g} = 0$.*

Proof. The proof is entirely analogous to the proof of Proposition 4.10. We indicate only the major steps. Set $U = U_{E^{\text{gen}},F^{\text{gen}}}$, and $U^\dagger = U_{E^{\text{gen}},F^{\text{gen}}}^\dagger$. Applying Proposition 1.2 to $V^\circ \rightarrow U^\dagger$ and $U^\dagger \rightarrow U$, we get

$$\text{rk Pic}_{\mathbf{Q}} U \leq 1 + \text{rk } \chi(G).$$

Let e be the number of divisorial components of $\tilde{\mathcal{H}}_{5,g} \setminus U$. We then get

$$\text{rk Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{5,g} \leq 1 + \text{rk } \chi(G) + e.$$

Both G and e depend on g modulo 4 and 5. Using Proposition 5.1 and Proposition 5.2, we get the following possibilities.

	$\text{rk } \chi(G) = \text{rk } \chi(G_E) + \text{rk } \chi(G_F)$	e
$4 \mid g, 5 \mid g + 4$	$0 = 0 + 0$	2 (M and CE)
$4 \mid g, 5 \nmid g + 4$	$1 = 0 + 1$	1 (M)
$4 \nmid g, 5 \mid g + 4$	$1 = 1 + 0$	1 (CE)
$4 \nmid g, 5 \nmid g + 4$	$2 = 1 + 1$	0 .

In all the cases, we have $\text{Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{5,g} \leq 3$. With Proposition 2.15, this gives $\text{Pic}_{\mathbf{Q}} \mathcal{H}_{5,g} = 0$. □

6 From Hurwitz spaces to Severi varieties

The associated scroll construction in § 2.1 lets us relate the Picard rank of a Hurwitz space to the Picard rank of a Severi variety. In this section, we work out this relation.

Recall the notation $\mathcal{U}_g(\mathbf{F}_m, d\tau)$, $\mathcal{V}_g(\mathbf{F}_m, d\tau)$, and $\mathcal{V}_g^{\text{irr}}(\mathbf{F}_m, d\tau)$ from § 0.1. When confusion is unlikely, we abbreviate them by \mathcal{U} , \mathcal{V} , and \mathcal{V}^{irr} . Following Diaz and Harris [DH88a], we enlarge \mathcal{U} by including the irreducible curves of geometric genus g having a cusp, a tacnode, a triple point, and irreducible nodal curves of geometric genus $(g - 1)$ (that is, curves having an “additional” node). Denote by $\tilde{\mathcal{U}}$ the normalization of this partial compactification. The local analysis from [DH88a, § 1] of the Severi variety at points corresponding to cusps, tacnodes, triple points, and an additional node shows that $\tilde{\mathcal{U}}$ is smooth. Since $\tilde{\mathcal{U}}$ maps to the linear series $|d\tau|$, it carries over it a family of (singular) curves. The normalization of the total space of this family gives a family $\mathcal{C} \rightarrow \tilde{\mathcal{U}}$ of curves

of arithmetic g . A generic fiber of $\mathcal{C} \rightarrow \tilde{\mathcal{U}}$ is the normalization the corresponding curve on \mathbf{F}_m .

Using the universal family, we can construct tautological divisor classes on $\tilde{\mathcal{U}}$ as follows. Consider the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\nu} & \mathbf{F}_m \\ \rho \downarrow & & \\ \tilde{\mathcal{U}} & & \end{array}$$

Define five tautological divisor classes on $\tilde{\mathcal{U}}$ (The subscript s stands for ‘‘Severi’’):

1. $\lambda_s := c_1(\rho_*\omega_\rho)$
2. $\kappa_s := \rho_*(c_1(\omega_\rho)^2)$
3. $\xi_s := \rho_*(\nu^*(f) \cdot c_1(\omega_\rho))$
4. $\theta_s := \rho_*(\nu^*(\sigma) \cdot c_1(\omega_\rho))$
5. $\psi_s := \rho_*(\nu^*[\text{Point}])$

Since the irreducible curves in the linear system $|d\tau|$ avoid the directrix σ , we get $\theta_s = \psi_s = 0$. Therefore, a natural conjecture is the following.

Conjecture 6.1. *The rational Picard group of $\tilde{\mathcal{U}}$ is tautological, that is*

$$\text{Pic}_{\mathbf{Q}}\tilde{\mathcal{U}} = \mathbf{Q}\langle \lambda_s, \kappa_s, \xi_s \rangle.$$

Denote by CU , TN , TP , and Δ the closures in \mathcal{V}^{irr} of the locus curves with a cusp, tacnode, triple point, or an additional node, respectively. Abusing notation, denote their preimages in $\tilde{\mathcal{U}}$ by the same letters.

Remark 6.2. It is not hard to check that the classes in $\text{Pic}_{\mathbf{Q}}\tilde{\mathcal{U}}$ of CU , TN , TP , and Δ can be expressed as \mathbf{Q} -linear combinations of λ_s , κ_s , and ξ_s and vice versa. Conjecture 6.1 is therefore equivalent to

$$\text{Pic}_{\mathbf{Q}}\mathcal{U} = 0.$$

Proposition 6.3. *The only divisorial components of $\mathcal{V}^{\text{irr}} \setminus \mathcal{U}$ are CU , TN , TP , and Δ .*

Proof. It suffices to show that the codimension one components of $\mathcal{V} \setminus \mathcal{U}$ are the loci of curves with cusps, tacnodes, triple points or an additional node. This follows by the same proof as for Theorem 1.4 in [DH88b]. The critical ingredient of the argument is provided by Lemma 6.4. □

Lemma 6.4. *Let $D \in |d\tau|$ be a reduced irreducible curve on the Hirzebruch surface \mathbf{F}_m . Denote by A the conductor ideal of the singularities of D . Then A imposes independent conditions on $H^0(\mathbf{F}_m, O(d\tau))$.*

Proof. Let $K = K_{\mathbf{F}_m}$ be the canonical class. The anti-canonical class $-K$ is effective. Furthermore, the fixed component of $-K$ is the directrix σ , and $-K$ separates points away from σ .

It is a classical that A imposes independent conditions on the adjoint linear system $|K + D|$. Let $Z = V(A)$ be the zero dimensional scheme defined by the ideal sheaf A . Then the restriction map

$$H^0(O(K + D)) \rightarrow H^0(O_Z(K + D))$$

is surjective. Therefore, we can conclude the same for

$$H^0(O(D)) \rightarrow H^0(O_Z(D))$$

by multiplying the previous restriction map by a general section of $O(-K)$. □

We now rephrase the Picard rank conjecture for Hurwitz spaces in a manner similar to Conjecture 6.1. Consider the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\alpha} & \mathbf{P}^1 \\ f \downarrow & & \\ \widetilde{\mathcal{H}}_{d,g}^\dagger & & \end{array}$$

Define the following tautological divisor classes on $\widetilde{\mathcal{H}}_{d,g}^\dagger$ (The subscript “h” stands for “Hurwitz”):

1. $\lambda_h := c_1(f_*\omega_f)$
2. $\kappa_h := f_*(c_1(\omega_f)^2)$
3. $\xi_h := f_*(\alpha^*[\text{Point}] \cdot c_1(\omega_f))$

Conjecture 6.5. *The rational Picard group of $\widetilde{\mathcal{H}}_{d,g}^\dagger$ is tautological, that is, $\text{Pic}_{\mathbf{Q}} \widetilde{\mathcal{H}}_{d,g}^\dagger = \mathbf{Q}\langle \lambda_h, \kappa_h, \xi_h \rangle$.*

Remark 6.6. It is easy to see that the classes of T , D , and Δ can be expressed as \mathbf{Q} -linear combinations of λ_h , κ_h , and ξ_h and vice versa. Also, by Proposition 1.3, the framed/unframed distinction is irrelevant. Therefore, Conjecture 6.5 is equivalent to the Picard rank conjecture stated in the introduction, namely that

$$\text{Pic}_{\mathbf{Q}} \mathcal{H}_{d,g} = 0.$$

We may thus drop the subscripts and use λ , κ , and ξ to denote the corresponding divisors on any of the spaces in (6.1).

Before we proceed, we must comment on the inclusion $X \hookrightarrow V$. The complement consists of $([\alpha: C \rightarrow \mathbf{P}^1], v)$, where $v \in H^0(C, \alpha^*O(m))$ does not give a birational map to \mathbf{F}_m . Let us disregard the α 's that factor non-trivially (such α 's form set of codimension at least two). Then the only such v are the pullbacks of the sections in $H^0(\mathbf{P}^1, O(m))$. The locus $([\alpha: C \rightarrow \mathbf{P}^1], v)$, where $v \in \alpha^*H^0(\mathbf{P}^1, O(m))$ has codimension at least two except in the case $g \equiv -1 \pmod{(d-1)}$, and $m = \lfloor (g+d-1)/(d-1) \rfloor$, that is, when the generic splitting of α_*O_C is

$$\alpha_*O_C = O \oplus O(-m) \oplus O(-m-1) \oplus \cdots \oplus O(-m-1).$$

In this case, the complement of X in V has a divisorial component given by the image of the constant vector bundle $H^0(\mathbf{P}^1, O(m)) \otimes O_W$. However, the class of this divisor in $\text{Pic}_{\mathbf{Q}} V \cong \text{Pic}_{\mathbf{Q}} W$ is in the span of λ , κ , and ξ . Therefore, in any case, $\text{Pic}_{\mathbf{Q}} V$ is spanned by λ , κ , and ξ if and only if $\text{Pic}_{\mathbf{Q}} X$ is.

Assume that Conjecture 6.1 holds. From diagram (6.1), we see that $\text{Pic}_{\mathbf{Q}} X$ is spanned by λ , κ , and ξ . By the comment about $X \hookrightarrow V$ above, this implies that $\text{Pic}_{\mathbf{Q}} V$, and in turn $\text{Pic}_{\mathbf{Q}} \tilde{\mathcal{H}}_{d,g}^\dagger$ is spanned by λ , κ , and ξ . Hence Conjecture 6.5 holds.

Assume that $m \geq \lceil 2(g+d-1)/(d-1) \rceil$ and Conjecture 6.5 holds. Then, by Proposition 2.6 the inclusion $Z \hookrightarrow \tilde{\mathcal{U}}$ is in fact an isomorphism. Again, diagram (6.1) shows that $\text{Pic}_{\mathbf{Q}} \tilde{\mathcal{U}}$ is spanned by λ , κ , and ξ . Hence Conjecture 6.1 holds. \square

Acknowledgments

We are grateful to our advisor Joe Harris for suggesting this problem and for providing constant support and encouragement. We also thank Gabriel Bujokas, Dawei Chen, Maksym Fedorchuk, and Ravi Vakil for helpful comments and conversations.

References

- [BV12] Michele Bolognesi and Angelo Vistoli, *Stacks of trigonal curves*, Trans. Amer. Math. Soc. **364** (2012), no. 7, 3365–3393. MR 2901217
- [Cas96] Gianfranco Casnati, *Covers of algebraic varieties. II. Covers of degree 5 and construction of surfaces*, J. Algebraic Geom. **5** (1996), no. 3, 461 – 477.
- [Cas98] ———, *Covers of algebraic varieties. III. The discriminant of a cover of degree 4 and the trigonal construction*, Trans. Amer. Math. Soc. **350** (1998), no. 4, 1359–1378.

- [CE96] Gianfranco Casnati and Torsten Ekedahl, *Covers of algebraic varieties. I. A general structure theorem, covers of degree 3, 4 and Enriques' surfaces*, J. Algebraic Geom. **5** (1996), 439–460.
- [CH88] Maurizio Cornalba and Joe Harris, *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves*, Ann. Sci. École Norm. Sup. (4) **21** (1988), no. 3, 455–475. MR 974412 (89j:14019)
- [Cop99] Marc Coppens, *Existence of pencils with prescribed scrollar invariants of some general type*, Osaka J. Math. **36** (1999), no. 4, 1049–1057. MR 1745644 (2001g:14006)
- [Cor07] Maurizio Cornalba, *The Picard group of the moduli stack of stable hyperelliptic curves*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **18** (2007), no. 1, 109–115. MR 2314467 (2008b:14017)
- [DE96] Steven Diaz and Dan Edidin, *Towards the homology of Hurwitz spaces*, J. Differential Geom. **43** (1996), no. 1, 66–98. MR MR1424420 (98e:14028)
- [DH88a] Steven Diaz and Joe Harris, *Geometry of the Severi variety*, Trans. Amer. Math. Soc. **309** (1988), no. 1, 1–34. MR 957060 (89i:14018)
- [DH88b] ———, *Ideals associated to deformations of singular plane curves*, Trans. Amer. Math. Soc. (1988).
- [Har83] John Harer, *The second homology group of the mapping class group of an orientable surface*, Invent. Math. **72** (1983), no. 2, 221–239. MR 700769 (84g:57006)
- [Har86] Joe Harris, *On the Severi problem*, Invent. Math. **84** (1986), no. 3, 445–461. MR 837522 (87f:14012)
- [KKV89] Friedrich Knop, Hanspeter Kraft, and Thierry Vust, *The Picard group of a G-variety*, Algebraische Transformationsgruppen und Invariantentheorie (1989).
- [MFK94] David Mumford, John Fogarty, and Frances Kirwan, *Geometric invariant theory*, third ed., Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], vol. 34, Springer-Verlag, Berlin, 1994. MR 1304906 (95m:14012)
- [Moc95] Shinichi Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci. **31** (1995), no. 3, 355–441. MR 1355945 (96j:14017)

- [Ohb97] Akira Ohbuchi, *On some numerical relations of d -gonal linear systems*, J. Math. Tokushima Univ. **31** (1997), 7–10. MR 1618986 (99b:14005)
- [Rec73] Sevin Recillas, *Maps between Hurwitz spaces*, Bol. Soc. Mat. Mexicana (2) **18** (1973), 59–63.
- [SF00] Zvezdelina E. Stankova-Frenkel, *Moduli of trigonal curves*, J. Algebraic Geom. **9** (2000), no. 4, 607–662. MR Mr1775309 (2001h:14031)
- [Tei80] Bernard Teissier, *Résolution simultanée, I*, Séminaire sur les Singularités des Surfaces, Lecture Notes in Mathematics, vol. 777, Springer, Berlin, 1980, pp. 71–81.
- [Tyo07] Ilya Tyomkin, *On Severi varieties on Hirzebruch surfaces*, Int. Math. Res. Not. IMRN (2007), no. 23, Art. ID rnm109, 31. MR 2380003 (2008m:14048)