

Solution 7


2. $f(x) = \sin x \Rightarrow f'(x) = \cos x$, so $f(\frac{\pi}{6}) = \frac{1}{2}$ and $f'(\frac{\pi}{6}) = \frac{1}{2}\sqrt{3}$. Thus,

$$L(x) = f(\frac{\pi}{6}) + f'(\frac{\pi}{6})(x - \frac{\pi}{6}) = \frac{1}{2} + \frac{1}{2}\sqrt{3}(x - \frac{\pi}{6}) = \frac{1}{2}\sqrt{3}x + \frac{1}{2} - \frac{1}{12}\sqrt{3}\pi.$$

6. $g(x) = \sqrt[3]{1+x} = (1+x)^{1/3} \Rightarrow g'(x) = \frac{1}{3}(1+x)^{-2/3}$, so $g(0) = 1$ and $g'(0) = \frac{1}{3}$. Therefore, $\sqrt[3]{1+x} = g(x) \approx g(0) + g'(0)(x-0) = 1 + \frac{1}{3}x$.

$$\text{So } \sqrt[3]{0.95} = \sqrt[3]{1+(-0.05)} \approx 1 + \frac{1}{3}(-0.05) = 0.98\bar{3},$$

$$\text{and } \sqrt[3]{1.1} = \sqrt[3]{1+0.1} \approx 1 + \frac{1}{3}(0.1) = 1.0\bar{3}.$$



24. To estimate $e^{-0.015}$, we'll find the linearization of $f(x) = e^x$ at $a = 0$. Since $f'(x) = e^x$, $f(0) = 1$, and $f'(0) = 1$, we have

$$L(x) = 1 + 1(x - 0) = x + 1. \text{ Thus, } e^x \approx x + 1 \text{ when } x \text{ is near } 0, \text{ so } e^{-0.015} \approx -0.015 + 1 = 0.985.$$

27. $y = f(x) = \tan x \Rightarrow dy = \sec^2 x dx$. When $x = 45^\circ$ and $dx = -1^\circ$,

$$dy = \sec^2 45^\circ (-\pi/180) = (\sqrt{2})^2 (-\pi/180) = -\pi/90, \text{ so } \tan 44^\circ = f(44^\circ) \approx f(45^\circ) + dy = 1 - \pi/90 \approx 0.965.$$

44. (a) $g'(x) = \sqrt{x^2+5} \Rightarrow g'(2) = \sqrt{9} = 3$. $g(1.95) \approx g(2) + g'(2)(1.95 - 2) = -4 + 3(-0.05) = -4.15$.

$$g(2.05) \approx g(2) + g'(2)(2.05 - 2) = -4 + 3(0.05) = -3.85.$$


(b) The formula $g'(x) = \sqrt{x^2+5}$ shows that $g'(x)$ is positive and increasing. This means that the slopes of the tangent lines are positive and the tangents are getting steeper. So the tangent lines lie *below* the graph of g . Hence, the estimates in part (a) are too small.

5. Absolute maximum value is $f(4) = 5$; there is no absolute minimum value; local maximum values are $f(4) = 5$ and $f(6) = 4$; local minimum values are $f(2) = 2$ and $f(1) = f(5) = 3$.

6. There is no absolute maximum value; absolute minimum value is $g(4) = 1$; local maximum values are $g(3) = 4$ and $g(6) = 3$; local minimum values are $g(2) = 2$ and $g(4) = 1$.

8. Absolute minimum at 1, absolute maximum at 5,

local maximum at 2, local minimum at 4



$$36. h(p) = \frac{p-1}{p^2+4} \Rightarrow h'(p) = \frac{(p^2+4)(1) - (p-1)(2p)}{(p^2+4)^2} = \frac{p^2+4-2p^2+2p}{(p^2+4)^2} = \frac{-p^2+2p+4}{(p^2+4)^2}.$$

$$h'(p) = 0 \Rightarrow p = \frac{-2 \pm \sqrt{4+16}}{-2} = 1 \pm \sqrt{5}. \text{ The critical numbers are } 1 \pm \sqrt{5}. [h'(p) \text{ exists for all real numbers.}]$$

$$40. g(\theta) = 4\theta - \tan \theta \Rightarrow g'(\theta) = 4 - \sec^2 \theta. \quad g'(\theta) = 0 \Rightarrow \sec^2 \theta = 4 \Rightarrow \sec \theta = \pm 2 \Rightarrow \cos \theta = \pm \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3} + 2n\pi, \frac{5\pi}{3} + 2n\pi, \frac{2\pi}{3} + 2n\pi, \text{ and } \frac{4\pi}{3} + 2n\pi \text{ are critical numbers.}$$

Note: The values of θ that make $g'(\theta)$ undefined are not in the domain of g .

$$60. f(x) = x - \ln x, \left[\frac{1}{2}, 2\right]. \quad f'(x) = 1 - \frac{1}{x} = \frac{x-1}{x}. \quad f'(x) = 0 \Rightarrow x = 1. \quad [\text{Note that } 0 \text{ is not in the domain of } f.]$$

$f\left(\frac{1}{2}\right) = \frac{1}{2} - \ln \frac{1}{2} \approx 1.19$, $f(1) = 1$, and $f(2) = 2 - \ln 2 \approx 1.31$. So $f(2) = 2 - \ln 2$ is the absolute maximum value and $f(1) = 1$ is the absolute minimum value.

1. (a) f is increasing on $(1, 3)$ and $(4, 6)$. (b) f is decreasing on $(0, 1)$ and $(3, 4)$.
 (c) f is concave upward on $(0, 2)$. (d) f is concave downward on $(2, 4)$ and $(4, 6)$.
 (e) The point of inflection is $(2, 3)$.

8. (a) f is increasing on the intervals where $f'(x) > 0$, namely, $(2, 4)$ and $(6, 9)$.
 (b) f has a local maximum where it changes from increasing to decreasing, that is, where f' changes from positive to negative (at $x = 4$). Similarly, where f' changes from negative to positive, f has a local minimum (at $x = 2$ and at $x = 6$).
 (c) When f' is increasing, its derivative f'' is positive and hence, f is concave upward. This happens on $(1, 3)$, $(5, 7)$, and $(8, 9)$. Similarly, f is concave downward when f' is decreasing—that is, on $(0, 1)$, $(3, 5)$, and $(7, 8)$.
 (d) f has inflection points at $x = 1, 3, 5, 7$, and 8 , since the direction of concavity changes at each of these values.

10. (a) $f(x) = 4x^3 + 3x^2 - 6x + 1 \Rightarrow f'(x) = 12x^2 + 6x - 6 = 6(2x^2 + x - 1) = 6(2x - 1)(x + 1)$. Thus,
 $f'(x) > 0 \Leftrightarrow x < -1$ or $x > \frac{1}{2}$ and $f'(x) < 0 \Leftrightarrow -1 < x < \frac{1}{2}$. So f is increasing on $(-\infty, -1)$ and $(\frac{1}{2}, \infty)$ and f is decreasing on $(-1, \frac{1}{2})$.

(b) f changes from increasing to decreasing at $x = -1$ and from decreasing to increasing at $x = \frac{1}{2}$. Thus, $f(-1) = 6$ is a local maximum value and $f\left(\frac{1}{2}\right) = -\frac{3}{4}$ is a local minimum value.

(c) $f''(x) = 24x + 6 = 6(4x + 1)$. $f''(x) > 0 \Leftrightarrow x > -\frac{1}{4}$ and $f''(x) < 0 \Leftrightarrow x < -\frac{1}{4}$. Thus, f is concave upward on $(-\frac{1}{4}, \infty)$ and concave downward on $(-\infty, -\frac{1}{4})$. There is an inflection point at $(-\frac{1}{4}, f(-\frac{1}{4})) = (-\frac{1}{4}, \frac{21}{8})$.

13. (a) $f(x) = \sin x + \cos x$, $0 \leq x \leq 2\pi$. $f'(x) = \cos x - \sin x = 0 \Rightarrow \cos x = \sin x \Rightarrow 1 = \frac{\sin x}{\cos x} \Rightarrow \tan x = 1 \Rightarrow x = \frac{\pi}{4}$ or $\frac{5\pi}{4}$. Thus, $f'(x) > 0 \Leftrightarrow \cos x - \sin x > 0 \Leftrightarrow \cos x > \sin x \Leftrightarrow 0 < x < \frac{\pi}{4}$ or $\frac{5\pi}{4} < x < 2\pi$ and $f'(x) < 0 \Leftrightarrow \cos x < \sin x \Leftrightarrow \frac{\pi}{4} < x < \frac{5\pi}{4}$. So f is increasing on $(0, \frac{\pi}{4})$ and $(\frac{5\pi}{4}, 2\pi)$ and f is decreasing on $(\frac{\pi}{4}, \frac{5\pi}{4})$.


(b) f changes from increasing to decreasing at $x = \frac{\pi}{4}$ and from decreasing to increasing at $x = \frac{5\pi}{4}$. Thus, $f(\frac{\pi}{4}) = \sqrt{2}$ is a local maximum value and $f(\frac{5\pi}{4}) = -\sqrt{2}$ is a local minimum value.

(c) $f''(x) = -\sin x - \cos x = 0 \Rightarrow -\sin x = \cos x \Rightarrow \tan x = -1 \Rightarrow x = \frac{3\pi}{4}$ or $\frac{7\pi}{4}$. Divide the interval $(0, 2\pi)$ into subintervals with these numbers as endpoints and complete a second derivative chart.

Interval	$f''(x) = -\sin x - \cos x$	Concavity
$(0, \frac{3\pi}{4})$	$f''(\frac{\pi}{2}) = -1 < 0$	downward
$(\frac{3\pi}{4}, \frac{7\pi}{4})$	$f''(\pi) = 1 > 0$	upward
$(\frac{7\pi}{4}, 2\pi)$	$f''(\frac{11\pi}{6}) = \frac{1}{2} - \frac{1}{2}\sqrt{3} < 0$	downward

There are inflection points at $(\frac{3\pi}{4}, 0)$ and $(\frac{7\pi}{4}, 0)$.


30. (a) $f(3) = 2 \Rightarrow$ the point $(3, 2)$ is on the graph of f . $f'(3) = \frac{1}{2} \Rightarrow$ the slope of the tangent line at $(3, 2)$ is $\frac{1}{2}$. $f'(x) > 0$ for all $x \Rightarrow f$ is increasing on \mathbb{R} . $f''(x) < 0$ for all $x \Rightarrow f$ is concave downward on \mathbb{R} . A possible graph for f is shown.



(b) The tangent line at $(3, 2)$ has equation $y - 2 = \frac{1}{2}(x - 3)$, or $y = \frac{1}{2}x + \frac{1}{2}$, and x -intercept -1 . Since f is concave downward on \mathbb{R} , f is below the x -axis at $x = -1$, and hence changes sign at least once. Since f is increasing on \mathbb{R} , it changes sign at most once. Thus, it changes sign exactly once and there is one solution of the equation $f(x) = 0$.

(c) $f'' < 0 \Rightarrow f'$ is decreasing. Since $f'(3) = \frac{1}{2}$, $f'(2)$ must be greater than $\frac{1}{2}$, so no, it is not possible that $f'(2) = \frac{1}{3}$.

32. (a) f is increasing where f' is positive, on $(1, 6)$ and $(8, \infty)$, and decreasing where f' is negative, on $(0, 1)$ and $(6, 8)$.
- (b) f has a local maximum where f' changes from positive to negative, at $x = 6$, and local minima where f' changes from negative to positive, at $x = 1$ and at $x = 8$.
- (c) f is concave upward where f' is increasing, that is, on $(0, 2)$, $(3, 5)$, and $(7, \infty)$, and concave downward where f' is decreasing, that is, on $(2, 3)$ and $(5, 7)$.
- (d) There are points of inflection where f changes its direction of concavity, at $x = 2$, $x = 3$, $x = 5$ and $x = 7$.



44. (a) $S(x) = x - \sin x$, $0 \leq x \leq 4\pi \Rightarrow S'(x) = 1 - \cos x$. $S'(x) = 0 \Leftrightarrow \cos x = 1 \Leftrightarrow x = 0, 2\pi$, and 4π .
 $S'(x) > 0 \Leftrightarrow \cos x < 1$, which is true for all x except integer multiples of 2π , so S is increasing on $(0, 4\pi)$ since $S'(2\pi) = 0$.

(b) There is no local maximum or minimum.

- (c) $S''(x) = \sin x$. $S''(x) > 0$ if $0 < x < \pi$ or $2\pi < x < 3\pi$, and $S''(x) < 0$ if $\pi < x < 2\pi$ or $3\pi < x < 4\pi$. So S is CU on $(0, \pi)$ and $(2\pi, 3\pi)$, and S is CD on $(\pi, 2\pi)$ and $(3\pi, 4\pi)$. There are inflection points at (π, π) , $(2\pi, 2\pi)$, and $(3\pi, 3\pi)$.

